


Article

Factors related to on-the-job abuse of nurses by patients

by Margot Shields and Kathryn Wilkins

April, 2009



Factors related to on-the-job abuse of nurses by patients

by Margot Shields and Kathryn Wilkins

Abstract

Background

Numerous studies indicate that health care providers, particularly nurses, face a high risk of on-the-job abuse from patients. This article examines physical and emotional abuse from patients in nurses working in hospitals or long-term care facilities.

Data and methods

Data are from the 2005 National Survey of the Work and Health of Nurses. Cross-tabulations were used to examine abuse in relation to personal characteristics of the nurse, job characteristics, and workplace climate factors. Multiple logistic regression modeling was used to examine abuse in relation to staffing and resource adequacy and relations among colleagues, controlling for personal and job characteristics.

Results

In 2005, 34% of Canadian nurses providing direct care in hospitals or long-term care facilities reported physical assault by a patient in the previous year; 47% reported emotional abuse. Abuse was related to being male, having less experience, usually working non-day shifts, and perceiving staffing or resources as inadequate, nurse-physician relations as poor, and co-worker and supervisor support as low. Associations between abuse and staffing or resource inadequacy and poor working relations persisted when controlling for personal and job characteristics.

Interpretation

Modifiable factors are important to nurses' on-the-job safety.

Keywords

resource allocation, violence, workload, workplace

Author

Margot Shields (613-951-4177; Margot.Shields@statcan.gc.ca) and Kathryn Wilkins (613-951-1769; Kathryn.Wilkins@statcan.gc.ca) are with the Health Information and Research Division at Statistics Canada, Ottawa, Ontario, K1A 0T6.

Health care providers are subject to a particularly high risk of workplace violence, and nurses are most at risk.¹⁻⁹

Evidence from numerous studies indicates that on-the-job abuse can result in a variety of negative outcomes among nurses, including anger, fear, depression, anxiety, sleep disruption, increased sick leave, symptoms of post-traumatic stress disorder, and job dissatisfaction.^{3,10-20} In addition, the likelihood of intending to leave their jobs or even the nursing profession altogether is greater among nurses who have experienced abuse on the job.^{14,20-24}

Abuse of health care providers may also affect recipients of care. Most of the research examining abuse in relation to quality of care is based on nurses' perceived ability to care for patients following incidents of abuse. Typical reports by nurses of the effects of abuse include impaired job performance, decreased productivity, and increased error.^{11,20-22,25}

The Canadian Nurses Association (CNA) and the International Council of Nurses (ICN) strongly advocate "zero-tolerance" of violence in the workplace.^{26,27} An improved understanding of the factors associated

with on-the-job abuse is an important prerequisite to the development of effective workplace policies.

Conceptual models of the factors that give rise to workplace violence in the health care sector generally include three levels of variables: individual characteristics of the nurse and the patient, workplace factors, and societal influences.²⁸ However, evidence-based research that considers variables from all three levels is scarce. A few studies²⁹⁻³² have been based on multivariate models that address both personal and workplace factors, but such research is relatively

uncommon, despite the view that it could inform the development of programs aimed at reducing on-the-job abuse of nurses.³³

This study examines the extent to which Canada's nurses working in hospitals or long-term care facilities face on-the-job abuse from patients. Then, on-the-job abuse is examined in relation to three groupings of variables—personal characteristics of the nurse, job characteristics, and workplace climate factors. A final objective is to determine whether workplace climate factors—of interest because of their potential for modification—are associated with abuse, independent of the potentially confounding effects of personal and job characteristics. The workplace climate factors studied are staffing and resource adequacy, nurse-physician working relations, and support from co-workers and supervisors.

Methods

Data source

The data for this study are from the 2005 National Survey of the Work and Health of Nurses (NSWHN), a comprehensive survey of employed regulated Canadian nurses (registered nurses, licensed practical nurses, and registered psychiatric nurses) conducted by Statistics Canada in partnership with the Canadian Institute for Health Information and Health Canada.³⁴ The purpose of the NSWHN was to collect information from nurses in all provinces and territories about their work environment, workload, perceived quality of patient care, and their physical and mental health. Survey content was included to provide data for analysis focusing on links between the nursing practice environment and various nurse and patient outcomes.

The NSWHN sample was selected at random from membership lists provided to Statistics Canada by all 26 provincial and territorial nursing organizations and regulating bodies across Canada. The survey was

administered by telephone over the period October 2005 to January 2006; the duration of a typical interview was 30 minutes.

Of the 24,443 nurses initially selected for the sample, 21,307 were successfully contacted, and of these, 1,015 were out-of-scope—meaning that they were not employed in nursing at the time of the survey. Another 1,616 (7.6% of the 21,307 who were contacted) refused to participate. Complete responses were obtained from 18,676 nurses, for a response rate of 79.7%.

Data were weighted to permit representative estimates of each of three nursing bodies—registered nurses (RNs), licensed practical nurses (LPNs) and registered psychiatric nurses (RPNs)—at the provincial level (and for the three territories combined).³⁴ Response rates by type of nurse were 80.8% among RNs, 78.4% among LPNs, and 80.6% among RPNs. Provincial response rates ranged from 77.0% in Ontario to 82.8% in Nova Scotia. The response rate in the three territories combined was 65.6%. The use of sampling weights is essential to reducing the potential for bias resulting from these differing response rates.

Reports of on-the-job abuse are far less common among nurses employed in settings such as community health, physicians' offices or educational institutions.³⁴ Therefore, to limit the heterogeneity of the sample, the analysis was restricted to nurses providing direct care to patients in hospitals or long-term care facilities. Of the total responding sample, 12,218 met these criteria; with survey weights applied, this sample was representative of the 218,300 Canadian nurses meeting the same criteria in the fall of 2005.

Definitions

Two yes/no questions were used to measure *on-the-job abuse* from patients:

- During the past 12 months, did you experience a physical assault from a patient?

- During the past 12 months, did you experience emotional abuse from a patient?

These two items were read to nurses; no further explanation or definition of physical assault or emotional abuse was given.

Four variables were used to assess *workplace climate*. Two of these variables, *staffing/resource adequacy* and *nurse-physician working relations*, are derived from subscales of the Nursing Work Index (NWI), a set of measures developed to study the nursing practice environment.³⁵ Response items were based on a four-point Likert scale: “strongly agree”—score 0, “somewhat agree”—score 1, “somewhat disagree”—score 2, “strongly disagree”—score 3.

The items comprising the staffing/resource adequacy subscale are shown in Table 1. A total staffing/resource adequacy score (with a possible range of 0 to 12) was calculated by summing the scores for the four items, with higher scores indicating lower levels of perceived adequacy. Cut-points were determined so as to divide the weighted distribution of scores into quartiles. In the NSWHN, the reliability coefficient (as assessed by the Cronbach's alpha) for this subscale was satisfactory, at 0.84, and satisfactory validity statistics have been previously reported.³⁶

Three statements measured nurse-physician working relations (Table 1); a total score (with a possible range of 0 to 9) was calculated by summing the scores for the three items; higher scores indicated less favourable relations. The weighted distribution of the scores was divided into quartiles. Cronbach's alpha for the nurse-physician working relations subscale was 0.82.

To maximize the number of respondents for whom scores were calculated, one “not applicable” or “not stated” response was accepted for both the staffing/resource adequacy and the nurse-physician working relations subscales. A score was calculated based on the items with responses and then adjusted by the

Table 1
Selected characteristics of nurses providing direct care in hospitals or long-term care facilities, Canada, 2005

	Sample size	Estimated number	Percent
Total	12,218	218,300	
Personal characteristics			
Female	11,365	205,400	94.1
Average years of experience in nursing (standard deviation)	17.0 (11.1)	17.1 (11.3)	...
Bachelor's degree or higher in nursing	1,653	42,200	19.4
Fair or poor general health	856	15,200	7.0
Fair or poor mental health	630	12,900	5.9
Job satisfaction			
Very satisfied	4,713	77,200	35.4
Somewhat satisfied	5,912	110,700	50.8
Somewhat dissatisfied	1,238	23,500	10.8
Very dissatisfied	328	6,400	2.9
Type of nurse			
Registered nurse	5,616	164,200	75.2
Licensed practical nurse	5,618	51,200	23.5
Registered psychiatric nurse	984	2,900	1.3
Job characteristics			
Work setting			
Hospital	8,081	172,100	78.8
Long-term care facility	4,137	46,200	21.2
Works full-time			
	6,938	127,000	58.4
Shift usually worked			
Days	3,370	68,600	31.4
Evenings	1,050	20,200	9.3
Nights	1,152	24,000	11.0
Mixed	6,644	105,400	48.3
Usually works 12-hour shift			
	4,453	75,600	36.8
Workplace climate factors			
Staffing/Resource adequacy			
Adequate support services allow me to spend time with my patients (percent disagreeing)	5,619	99,900	47.0
There is enough time and opportunity to discuss patient care (percent disagreeing)	4,602	91,800	42.5
There are enough nurses on staff to provide quality patient care (percent disagreeing)	6,403	121,000	55.9
There is enough staff to get the work done (percent disagreeing)	5,854	112,000	51.7
Nurse-physician working relations			
There is a lot of teamwork between nurses and physicians (percent disagreeing)	2,310	41,100	19.2
There is collaboration between nurses and physicians (percent disagreeing)	1,271	24,300	11.3
Physicians and nurses have good working relations (percent disagreeing)	1,475	28,300	13.1
Supervisor support			
Your supervisor is helpful in getting the job done (percent disagreeing)	3,267	59,700	27.7
Co-worker support			
You are exposed to hostility or conflict from the people you work with (percent agreeing)	5,508	100,200	46.2
The people you work with are helpful in getting the job done (percent disagreeing)	413	8,200	3.8

... not applicable

Note: Because of missing values, percent may not correspond to estimated number divided by total.

Source: 2005 National Survey of the Work and Health of Nurses.

mean substitution technique to compensate for the item without a response.

Two statements were used to measure *co-worker support*:

- You were exposed to hostility or conflict from the people you work with.
- The people you work with were helpful in getting the job done.

Response options were: “strongly agree,” “agree,” “neither agree nor disagree,” “disagree,” or “strongly disagree.” Respondents were classified as having low co-worker support if they indicated “strongly agree” or “agree” in response to the first item, or “disagree” or “strongly disagree” in response to the second.

Supervisor support was measured with the item, “Your supervisor is helpful in getting the job done.” Respondents were classified as having low supervisor support if they indicated “disagree” or “strongly disagree.”

Detailed definitions and questionnaire items for the personal and job characteristics used in this study are available in a previously published report.³⁴

Analytical techniques

In processing the NSWHN data, Statistics Canada methodologists produced survey weights so that the data were representative of all regulated nurses across Canada. This analysis is based on data weighted to be representative of nurses employed in hospitals or long-term care facilities who provide direct care to patients. Frequency estimates were produced to examine characteristics of the study population. Bivariate estimates were used to examine factors associated with physical assault and emotional abuse from patients among these nurses.

Logistic regression models were used to examine abuse in relation to workplace climate factors. Three sets of models were fitted. In the first set, unadjusted odds were calculated to examine the individual relationship of each workplace climate factor to abuse. In the second set, personal characteristics of the nurse and job characteristics were included as control variables. Among the control variables reflecting personal characteristics, attitudinal factors were considered important—in particular, a generally gloomy outlook—because of the possible influence on the perception or likelihood of a nurse’s reporting on-the-job abuse.

In the absence of variables directly measuring negative affectivity, self-reports of poor mental health and job dissatisfaction were used as control variables. The final model included all four workplace climate factors, in addition to personal and job characteristics. This was done to determine if workplace climate factors—of interest because of their potential to be changed—were associated with abuse, independent of the potentially confounding effects of personal and job characteristics. Selection of the personal and job characteristic control variables was guided by findings in the literature and availability in the NSWHN.

The bootstrap technique³⁷ was used to estimate standard errors, coefficients of variation and 95% confidence intervals. Differences between estimates were tested for statistical significance established at the level of $p < 0.05$.

Results

Characteristics of study population

In 2005, the number of nurses delivering direct patient care in hospitals or long-term care facilities was estimated at just over 218,000, based on a weighted sample of 12,218 respondents (Table 1). The overwhelming majority (94%) were women. On average, they had 17 years of experience as a nurse. Just under one-fifth (19%) had a bachelor's degree or higher in nursing. Most were in good health; only 7% rated their general health as "fair" or "poor," and 6% rated their mental health as "fair" or "poor." The vast majority were satisfied with their jobs—35% were very satisfied and 51%, somewhat satisfied. Three-quarters were registered nurses (RNs); 24% were licensed practical nurses (LPNs); and the remaining 1% were registered psychiatric nurses (RPNs).

Over half (58%) of nurses in the study population were employed full-time, and close to four-fifths (79%) worked in hospitals. Two-thirds (69%) worked shifts other than exclusively

Table 2
Number and percentage reporting physical assault by a patient over past 12 months, by selected characteristics, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005

	Estimated number	Percent	95% confidence interval	
			from	to
Total	73,300	33.8	32.5	35.1
Personal characteristics				
Sex				
Female	67,400	33.0*	31.7	34.4
Male [†]	5,900	46.1	40.6	51.6
Years of experience in nursing				
Fewer than 5	16,300	41.7*	38.7	44.7
5 to 9	10,500	37.6	33.8	41.5
10 to 14	10,300	36.9	33.3	40.5
15 to 19 [†]	10,600	33.8	30.3	37.3
20 to 24	8,700	32.7	29.3	36.2
25 to 29	9,500	27.8*	24.9	30.6
30 or more	7,400	24.9*	21.6	28.1
Bachelor's degree or higher in nursing				
Yes	12,400	29.4*	26.4	32.5
No [†]	60,900	34.8	33.4	36.3
General health				
Good, very good or excellent	67,600	33.5	32.1	34.8
Fair or poor [†]	5,700	38.2	33.2	43.2
Mental health				
Good, very good or excellent	67,600	33.1*	31.8	34.5
Fair or poor [†]	5,700	45.2	39.3	51.2
Job satisfaction				
Very satisfied [†]	21,100	27.4	25.4	29.5
Somewhat satisfied	37,900	34.5*	32.6	36.3
Somewhat dissatisfied	10,800	46.5*	42.1	50.9
Very dissatisfied	3,300	53.1*	44.7	61.5
Type of nurse				
Registered nurse [†]	49,100	30.2	28.5	31.8
Licensed practical nurse	22,800	44.8*	43.0	46.5
Registered psychiatric nurse	1,400	47.2*	44.2	50.1
Job characteristics				
Work setting				
Hospital	50,500	29.6*	28.0	31.1
Long-term care facility [†]	22,800	49.6	47.3	51.8
Work status				
Full-time	43,400	34.4	32.7	36.2
Part-time [†]	29,600	32.9	30.9	34.9
Shift usually worked				
Days [†]	15,900	23.3	21.2	25.4
Evenings	8,100	40.2*	35.8	44.5
Nights	9,200	38.7*	34.5	43.0
Mixed	40,100	38.3*	36.4	40.2
Length of shift				
12 hours	28,800	38.5*	36.0	40.9
Under 12 hours [†]	40,000	31.0	29.5	32.6
Workplace climate factors				
Staffing/Resource adequacy				
First quartile (most adequate)	9,800	23.4	21.0	25.9
Second quartile	14,400	28.7 [†]	26.1	31.3
Third quartile	22,500	35.3 [†]	32.9	37.8
Fourth quartile (least adequate)	26,100	44.1 [†]	41.4	46.7
Nurse-physician working relations				
First quartile (most favourable)	14,400	28.2	25.7	30.8
Second quartile	19,300	33.7 [†]	31.1	36.3
Third quartile	20,100	34.7	32.4	37.0
Fourth quartile (least favourable)	18,300	38.9 [†]	36.1	41.6
Low supervisor support				
Yes	23,700	39.8*	37.2	42.5
No [†]	49,200	31.7	30.2	33.1
Low co-worker support				
Yes	40,300	39.7*	37.7	41.7
No [†]	32,900	28.7	27.1	30.4

[†] reference category

* significantly different from estimate for reference category ($p < 0.05$)

[†] significantly different from estimate for previous quartile ($p < 0.05$)

Source: 2005 National Survey of the Work and Health of Nurses.

days, and 37% reported usually working a 12-hour shift.

Workplace climate

Substantial percentages of nurses in the study population perceived that staffing or resources were less than adequate. The majority disagreed that there were enough nurses on staff to provide quality patient care (56%), or enough to get the work done (52%). Slightly lower percentages disagreed that adequate support services allowed them time to spend with patients (47%), and that there was enough time and opportunity to discuss patient care (43%).

In contrast, problems regarding relations with physicians were reported infrequently. A lack of teamwork between nurses and physicians was reported by 19%, and a lack of collaboration, by 11%. Thirteen percent disagreed with the statement “Physicians and nurses have good working relations.”

Just over one-quarter (28%) reported that their supervisor was not helpful in getting the job done.

Although very few (4%) disagreed that the people they worked with were helpful in getting the job done, close to half (46%) reported that they were exposed to hostility or conflict from co-workers.

Factors associated with abuse

Among nurses working in hospitals or long-term care facilities, 34% reported physical assault from a patient over the past year (Table 2), and 47% reported emotional abuse (Table 3). Male nurses and less experienced nurses were more likely to report both types of abuse. Having a bachelor’s degree or higher in nursing was associated with a decreased likelihood of reporting physical assault, but was not related to emotional abuse. Compared with RNs, LPNs and RPNs were more likely to report abuse. RPNs were particularly at risk, with 47% reporting physical assault and 72% reporting emotional abuse.

Table 3
Number and percentage reporting emotional abuse by a patient over past 12 months, by selected characteristics, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005

	Estimated number	Percent	95% confidence interval	
			from	to
Total	101,200	46.7	45.3	48.1
Personal characteristics				
Sex				
Female	94,200	46.2*	44.8	47.6
Male [†]	7,000	54.6	48.7	60.4
Years of experience in nursing				
Fewer than 5	18,800	48.2	44.9	51.5
5 to 9	13,600	48.7	44.8	52.7
10 to 14	14,600	52.1	48.4	55.9
15 to 19 [†]	16,300	51.8	48.1	55.5
20 to 24	12,300	46.6	42.9	50.4
25 to 29	14,000	40.8*	37.3	44.3
30 or more	11,500	39.1*	35.2	42.9
Bachelor's degree or higher in nursing				
Yes	19,300	46.1	42.6	49.5
No [†]	81,800	46.8	45.3	48.4
General health				
Good, very good or excellent	92,900	46.0*	44.6	47.5
Fair or poor [†]	8,300	56.0	50.8	61.3
Mental health				
Good, very good or excellent	93,100	45.6*	44.2	47.0
Fair or poor [†]	8,000	64.3	58.4	70.3
Job satisfaction				
Very satisfied [†]	29,700	38.7	36.4	41.0
Somewhat satisfied	53,500	48.6*	46.6	50.6
Somewhat dissatisfied	13,400	57.7*	53.4	62.0
Very dissatisfied	4,200	68.5*	60.9	76.1
Type of nurse				
Registered nurse [†]	74,400	45.7	43.9	47.5
Licensed practical nurse	24,700	48.6*	46.8	50.3
Registered psychiatric nurse	2,100	71.6*	69.0	74.2
Job characteristics				
Work setting				
Hospital	79,100	46.3	44.7	48.0
Long-term care facility [†]	22,000	48.0	45.7	50.2
Work status				
Full-time	60,200	47.8	45.9	49.7
Part-time [†]	40,700	45.3	43.3	47.3
Shift usually worked				
Days [†]	25,000	36.6	34.2	39.1
Evenings	9,800	48.8*	44.6	53.0
Nights	11,400	47.9*	43.4	52.3
Mixed	55,000	52.6*	50.6	54.6
Length of shift				
12 hours	41,000	54.7*	52.1	57.3
Under 12 hours [†]	53,700	41.7	40.0	43.4
Workplace climate factors				
Staffing/Resource adequacy				
First quartile (most adequate)	13,200	31.7	28.7	34.7
Second quartile	20,800	41.4 [†]	38.6	44.2
Third quartile	32,100	50.5 [†]	47.9	53.0
Fourth quartile (least adequate)	34,400	58.0 [†]	55.4	60.6
Nurse-physician working relations				
First quartile (most favourable)	19,700	38.7	35.8	41.5
Second quartile	24,900	43.5 [†]	40.8	46.2
Third quartile	28,100	48.6 [†]	46.0	51.1
Fourth quartile (least favourable)	27,000	57.4 [†]	54.6	60.2
Low supervisor support				
Yes	32,500	54.6*	52.0	57.2
No [†]	68,200	43.9	42.3	45.6
Low co-worker support				
Yes	53,400	52.6*	50.7	54.6
No [†]	47,600	41.6	39.7	43.5

[†] reference category

* significantly different from estimate for reference category (p < 0.05)

[†] significantly different from estimate for previous quartile (p < 0.05)

Source: 2005 National Survey of the Work and Health of Nurses.

Nurses working shifts other than days and those who usually worked a 12-hour shift were more likely to report both types of abuse.

Reports of abuse varied substantially by clinical area of practice. The percentage of nurses reporting physical assault was particularly high among

those working in geriatrics/long-term care (50%), palliative care (47%), psychiatry/mental health (44%), critical care (44%), or the emergency room (42%) (Figure 1). Emotional abuse was more common among nurses working in psychiatry/mental health

(70%), the emergency room (69%), critical care (54%), medicine/surgery (52%) or geriatrics/long-term care (49%) (Figure 2).

Workplace climate and abuse


The four workplace climate factors considered in this study—staffing/resource adequacy, nurse-physician working relations, supervisor support, and co-worker support—were all significantly associated with both physical assault (Table 2) and emotional abuse (Table 3). The data were suggestive of a gradient between the risk of abuse and staffing/resource adequacy. Reports of physical assault were highest (44%) among nurses who perceived staffing or resources to be the least adequate (quartile 4) and lowest (23%) among those who thought they were the most adequate (quartile 1). The corresponding estimates for emotional abuse were 58% for quartile 4 and 32% for quartile 1.

A gradient was also observed between abuse and nurse-physician working relations. The percentage reporting physical assault ranged from 28% of nurses perceiving the most favourable relations to 39% of those perceiving relations as least favourable. A more pronounced gradient was observed for reports of emotional abuse: 39% of those in the most favourable quartile versus 57% of those in the least favourable.

Nurses classified as having low supervisor support were more likely to report physical assault, compared with those reporting more positive relations (40% versus 32%). The same was true for emotional abuse (55% versus 44%).

Similar differences emerged according to level of co-worker support; 40% of nurses with low co-worker support reported physical assault, compared with 29% of those with more supportive co-workers. For emotional abuse, the comparable figures were 53% versus 42%.

Figure 1
Percentage reporting physical assault by a patient over past 12 months, by clinical area of employment, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005




* significantly different from estimate for total

^E interpret with caution (coefficient of variation between 16.6% and 33.3%)

Source: 2005 National Survey of the Work and Health of Nurses.

Figure 2
Percentage reporting emotional abuse by a patient over past 12 months, by clinical area of employment, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005



* significantly different from estimate for total

Source: 2005 National Survey of the Work and Health of Nurses.

Multivariate analysis

Workplace climate factors were examined individually in multivariate models controlling for the influences of personal characteristics of the nurse and job characteristics (Table 4). Although controlling for these

potentially confounding variables somewhat reduced the strength of the associations, all four workplace climate factors remained significantly related to both physical assault and emotional abuse. Simultaneously including all four of the workplace climate factors

and personal and job characteristics further weakened the strength of associations of workplace climate factors with abuse, because of the correlations among the workplace climate factors. Nevertheless, perceptions that staffing or resources were inadequate or that

Table 4
Odds ratios relating workplace climate factors to physical assault/emotional abuse by a patient over past 12 months, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005

Workplace climate factors	Unadjusted odds ratio	95% confidence interval		Adjusted for personal and job characteristics			Adjusted for workplace climate, personal and job characteristics		
		from	to	Adjusted odds ratio	95% confidence interval from	to	Adjusted odds ratio	95% confidence interval from	to
Physical assault by patient									
Staffing/Resource adequacy									
First quartile (most adequate) [†]	1.0	1.0	1.0
Second quartile	1.3*	1.1	1.6	1.3*	1.0	1.6	1.2	1.0	1.5
Third quartile	1.8*	1.5	2.1	1.6*	1.3	2.0	1.5*	1.2	1.8
Fourth quartile (least adequate)	2.6*	2.2	3.1	2.3*	1.9	2.8	2.1*	1.7	2.6
Nurse-physician working relations									
First quartile (most favourable) [†]	1.0	1.0	1.0
Second quartile	1.3*	1.1	1.5	1.2	1.0	1.4	1.1	0.9	1.4
Third quartile	1.3*	1.1	1.6	1.2*	1.0	1.4	1.1	0.9	1.3
Fourth quartile (least favourable)	1.6*	1.4	1.9	1.3*	1.0	1.5	1.0	0.8	1.2
Low supervisor support									
Yes	1.4*	1.3	1.6	1.3*	1.2	1.6	1.2*	1.0	1.4
No [†]	1.0	1.0	1.0
Low co-worker support									
Yes	1.6*	1.5	1.8	1.6*	1.4	1.8	1.4*	1.3	1.7
No [†]	1.0	1.0	1.0
Emotional abuse from patient									
Staffing/Resource adequacy									
First quartile (most adequate) [†]	1.0	1.0	1.0
Second quartile	1.5*	1.3	1.8	1.4*	1.2	1.8	1.3*	1.1	1.6
Third quartile	2.2*	1.8	2.6	1.9*	1.6	2.4	1.7*	1.4	2.1
Fourth quartile (least adequate)	3.0*	2.5	3.5	2.6*	2.1	3.1	2.2*	1.8	2.6
Nurse-physician working relations									
First quartile (most favourable) [†]	1.0	1.0	1.0
Second quartile	1.2*	1.0	1.4	1.2*	1.0	1.4	1.1	0.9	1.3
Third quartile	1.5*	1.3	1.8	1.4*	1.2	1.7	1.3*	1.1	1.5
Fourth quartile (least favourable)	2.1*	1.8	2.5	1.9*	1.6	2.3	1.5*	1.3	1.8
Low supervisor support									
Yes	1.5*	1.4	1.7	1.4*	1.2	1.6	1.2*	1.0	1.4
No [†]	1.0	1.0	1.0
Low co-worker support									
Yes	1.6*	1.4	1.8	1.5*	1.3	1.7	1.3*	1.2	1.5
No [†]	1.0	1.0	1.0

[†] reference category

* significantly different from estimate for reference category (p < 0.05)

... not applicable

Note: Personal characteristics included in the models were sex, years of experience in nursing, nursing education, general health, mental health, job satisfaction, and type of nurse. Job characteristics included work setting, clinical area of employment, work status, shift usually worked, and length of shift. See Appendix Tables A and B for results of full models.

Source: 2005 National Survey of the Work and Health of Nurses.

co-worker or supervisor support were low remained positively associated with both types of abuse. Unfavourable relations among nurses and physicians remained positively associated with emotional abuse, but the association with physical assault did not persist.

Discussion

This study, based on data from a large nationally representative sample with an exceptionally high response rate, found that on-the-job abuse by patients is common among Canada's nurses. Studies elsewhere have also found that nurses face a high risk of on-the-job abuse, but that they tend to accept it as "part of the job".^{5-8,14,38-40} Many nurses do not bother to document incidents of violence, either because they feel that no action will be taken or that they will be held accountable; a "culture of silence" is said to exist.^{3,6,7,14,41-44}

Because of differing definitions of abuse, it is difficult to compare estimates from the NSWHN with those from other studies. A notable exception is the 2005 National Health Services (NHS) staff survey in England, in which nurses were asked questions similar to those used in the NSWHN: "In the past 12 months have you experienced physical violence from any of the following? Patients/Service users" and "In the past 12 months have you experienced harassment, bullying or abuse from any of the following? Patients/Service users."⁴⁵ Sixteen percent of NHS nurses reported physical abuse from a patient/service user over the past year, and 26% reported harassment, bullying or abuse. The NHS estimates are based on all nurses, regardless of work setting or job tasks. When estimates from the NSWHN were tabulated so that they were based on all nurses, 25% reported physical assault, and 38%, emotional abuse. Although comparisons with NHS estimates need to be interpreted with caution because of the somewhat different wording of the questions

(particularly for emotional abuse), the Canadian estimates are substantially higher than those from England.

A possible explanation for the lower rates among British nurses relates to support for reporting abuse and follow-through of such reports by authorities. Among the NHS nurses who experienced physical abuse, 69% indicated that they reported the incident; among those who experienced emotional abuse, 57% reported the incident. These figures are appreciably higher than estimates from other studies. For example, in a survey in 1998/1999 of RNs in hospitals in Alberta and British Columbia, only 36% of those who experienced physical abuse indicated that they had reported the incident to the hospital, and of those who experienced emotional abuse, 28%.³¹ A particularly relevant finding from the NHS survey was that very few nurses reported a lack of "effective action" when staff were either physically or emotionally abused. Encouragement for reporting incidents of abuse, along with an appropriate response by those in authority, may be required to reduce on-the-job abuse among nurses.

Consistent with other research, estimates from the NSWHN show that nurses with fewer years' experience^{13,21,29-31,46-48} and male nurses^{14,21,30,46} were more likely to report both physical and emotional abuse from patients. Nurses with less experience may not have the necessary skills to predict and defuse abusive situations. Alternatively, inexperienced, younger nurses may more readily acknowledge incidents of abuse, since they are less likely to accept it as being "part of the job."³¹ Reasons that have been proposed for the higher risk of abuse among male nurses include greater exposure to violent patients, and societal norms that differ between the sexes.^{14,21,48} One study found a tendency for male nurses to feel protective of female staff and to assume the primary role in restraining aggressive patients.⁴⁰

Job characteristics associated with reports of abuse in this study were shift work and clinical area of employment. Shift work—particularly the night shift—has been linked to abuse in other research,³⁰ and may be related to working in more isolated conditions. Also consistent with other studies was the finding that nurses who work predominantly in psychiatry, emergency, geriatrics or long-term care, or critical care are particularly subject to abuse.^{14,15,21,29,30,32,33}

An important finding from this study was that perceiving that staffing or resources were inadequate was associated with both physical and emotional abuse, independent of the potentially confounding effects of personal and job characteristics. Although studies examining workplace climate factors in relation to on-the-job abuse are relatively scarce, somewhat similar results have emerged from other research. A survey of nursing staff from eight European countries found that time pressure, defined as the extent to which nurses lack time to accomplish tasks, was associated with on-the-job abuse.³⁰ Another study of RNs working in hospitals in Alberta and British Columbia found that nurses who reported they had left tasks undone in their last shift because of lack of time were more likely to report abuse.³¹ If nurses lack the time to complete necessary tasks as a result of staffing and resource inadequacy, patients may become agitated, thereby increasing the risk of violence directed at the nurse.

In this study, interpersonal relations were also related to abuse. Nurses who reported poor working relations with physicians, low supervisor support or low co-worker support were more likely to report abuse from patients. It has been hypothesized that hostile interactions among health care workers result in increased distress levels. In turn, relations between patients and nurses may be jeopardized.³⁰

What is already known on this subject?

- Health care providers commonly experience violence or verbal abuse from patients in their care, and nurses are particularly at risk.
- Nurses who experience on-the-job abuse are at risk of physical and psychological problems.
- There is also some evidence of a link between on-the-job abuse of nurses and diminished quality of patient care.

What does this study add?

- This is the first Canadian study based on nationally representative data to quantify the extent to which nurses working in hospitals or long-term care facilities report on-the-job abuse from patients, and to examine factors associated with abuse.
- Workplace climate factors—staffing and resource adequacy, relations between nurses and physicians, co-worker support and supervisor support—are negatively related to on-the-job abuse.
- Associations between workplace climate and abuse are independent of the effects of personal and job characteristics.

Limitations

Estimating the extent to which nurses experience on-the-job abuse and comparing estimates across surveys is hampered by the lack of a consistent definition of workplace violence.⁴⁹ Similar to other research, estimates of physical assault and emotional abuse

for this study were based on self-reported data from nurses. No further explanation or definitions of these terms were given to respondents, and estimates of abuse were not validated against more objective sources.

The one-year period over which abuse was measured may have resulted in recall bias. As well, the survey asked no questions about the frequency or the severity of the abuse, which would have made it possible to gain a more complete understanding of predictors of abuse.

Negative affectivity, or a general tendency to be pessimistic, may have influenced the likelihood of negative perceptions of workplace climate factors and reporting abuse. If so, exaggerated associations between workplace climate factors and abuse may have resulted. Including job satisfaction and perceived mental health as control variables may have partly addressed this limitation, depending on the extent to which negative affectivity is correlated with job satisfaction and mental health.

The associations observed in the analysis may have been partially accounted for by societal factors that could not be considered because of the unavailability of data from the NSWHN. For example, influences arising from the socio-political context, the economy or the geographic location of the health care facility may have affected the likelihood of reports of abuse, but measures of such factors were not available.

The measures of workplace climate factors in the NSWHN are based on reports from nurses. Different results may have emerged if more objective measures—such as nurse-patient ratios and professional staffing mix (the ratio of registered nurses to licensed practical nurses and auxiliary staff)—had been used. The design of the NSWHN precluded linkage to administrative data that contain this information.

The NSWHN data are cross-sectional, so the temporal ordering of factors observed to be associated with each other cannot be established,

and causality cannot be inferred. For example, whether nurses in fair or poor mental health were more likely than those in better mental health to be subsequently abused, or whether nurses who were abused were then more likely to experience fair or poor mental health, cannot be discerned from the data.

The NSWHN was administered to respondents by telephone. The degree to which this method of data collection may have affected the accuracy of responses is unknown.

Conclusion

Findings from the NSWHN reveal that a substantial proportion of Canada's nurses experience physical and emotional abuse at the hands of patients. Workplace climate factors, including the perception that staffing and resources are inadequate and that interpersonal relations among health care workers are poor, were found to be related to higher risks of on-the-job abuse from patients. The importance of these findings is underscored by numerous studies that have found associations between on-the-job abuse from patients and a host of physical and psychological problems among nurses. Furthermore, studies providing evidence of a link between abuse from patients and nursing caregiving errors suggest that nurses' role may be compromised as a consequence of abuse. These potentially harmful consequences and the pervasiveness of abuse of Canada's nurses emphasize the importance of staffing and resource adequacy and interpersonal relations among health care providers. ■

References

1. Aiken LH, Clarke SP, Sloane DM, et al. Nurses' reports on hospital care in five countries. *Health Affairs* 2001; 20(3): 43-53.
2. Chapman R, Styles I. An epidemic of abuse and violence: nurse on the front line. *Accident and Emergency Nursing* 2006; 14(4): 245-9.
3. Di Martino V. *Workplace Violence in the Health Sector. Country Case Studies*. Geneva: International Labour Office, International Council of Nurses, World Health Organization and Public Services International, 2002. Available at: <http://www.icn.ch/SynthesisReportWorkplaceViolenceHealthSector.pdf>. Accessed January 17, 2008
4. Elliott PP. Violence in health care. What nurse managers need to know. *Nursing Management* 1997; 28(12): 38-41.
5. Hewitt JB, Levin PF. Violence in the workplace. *Annual Review of Nursing Research* 1997; 15: 81-99.
6. International Council of Nurses. *Violence: A World-wide Epidemic (Fact Sheet)* Geneva: International Council of Nurses, 2004. Available at: http://www.icn.ch/matters_violence.htm. Accessed January 17, 2008.
7. Jones J, Lyneham J. Violence: part of the job for Australian nurses? *Australian Journal of Advanced Nursing* 2000; 18(2): 27-32.
8. McPhaul KM, Lipscomb JA. Workplace violence in health care: recognized but not regulated. *Online Journal of Issues in Nursing* 2004; 9(3): 7.
9. Yassi A, Gilbert M, Cvitkovich Y. Trends in injuries, illnesses, and policies in Canadian healthcare workplaces. *Canadian Journal of Public Health* 2005; 96(5): 333-9.
10. Arnetz JE, Arnetz BB. Violence towards health care staff and possible effects on the quality of patient care. *Social Science and Medicine* 2001; 52(3): 417-27.
11. Celik SS, Celik Y, Agirbas I, et al. Verbal and physical abuse against nurses in Turkey. *International Nursing Review* 2007; 54(4): 359-66.
12. Dougherty LM, Bolger JP, Preston DG, et al. Effects of exposure to aggressive behavior on job satisfaction of health care staff. *Journal of Applied Gerontology* 1992; 11(2): 160-72.
13. Evers W, Tomic W, Brouwers A. Aggressive behaviour and burnout among staff of homes for the elderly. *International Journal of Mental Health Nursing* 2002; 11(1): 2-9.
14. Gerberich SG, Church TR, McGovern PM, et al. An epidemiological study of the magnitude and consequences of work related violence: the Minnesota Nurses' Study. *Occupational and Environmental Medicine* 2004; 61(6): 495-503.
15. Hesketh KL, Duncan SM, Estabrooks CA, et al. Workplace violence in Alberta and British Columbia hospitals. *Health Policy* 2003; 63(3): 311-21.
16. Inoue M, Tsukano K, Muraoka M, et al. Psychological impact of verbal abuse and violence by patients on nurses working in psychiatric departments. *Psychiatry and Clinical Neurosciences* 2006; 60(1): 29-36.
17. Kivimaki M, Elovainio M, Vahtera J. Workplace bullying and sickness absence in hospital staff. *Occupational and Environmental Medicine* 2000; 57(10): 656-60.
18. Kivimaki M, Virtanen M, Vartiainen M, et al. Workplace bullying and the risk of cardiovascular disease and depression. *Occupational and Environmental Medicine* 2003; 60(10): 779-83.
19. O'Connell B, Young J, Brooks J, et al. Nurses' perceptions of the nature and frequency of aggression in general ward settings and high dependency areas. *Journal of Clinical Nursing* 2000; 9(4): 602-10.
20. Sofield L, Salmond SW. Workplace violence. A focus on verbal abuse and intent to leave the organization. *Orthopedic Nursing* 2003; 22(4): 274-83.
21. Farrell GA, Bobrowski C, Bobrowski P. Scoping workplace aggression in nursing: findings from an Australian study. *Journal of Advanced Nursing* 2006; 55(6): 778-87.
22. Fernandes CM, Bouthillette F, Raboud JM, et al. Violence in the emergency department: a survey of health care workers. *Canadian Medical Association Journal* 1999; 161(10): 1245-8.
23. Lanza ML, Zeiss R, Rierdan J. Violence against psychiatric nurses: sensitive research as science and intervention. *Contemporary Nurse* 2006; 21(1): 71-84.
24. Quine L. Workplace bullying in NHS community trust: staff questionnaire survey. *British Medical Journal* 1999; 318(7178): 228-32.
25. Rowe MM, Sherlock H. Stress and verbal abuse in nursing: do burned out nurses eat their young? *Journal of Nursing Management* 2005; 13(3): 242-8.
26. Canadian Nurses Association. *Violence in the Workplace (Fact Sheet)* Ottawa: Canadian Nurses Association, 2005. Available at: http://www.cna-nurses.ca/CNA/documents/pdf/publications/FS22_Violence_Workplace_e.pdf. Accessed January 17, 2008.
27. International Council of Nurses. *Abuse and Violence Against Nursing Personnel (Position Statement)* Geneva: International Council of Nurses, 2000. Available at: <http://www.icn.ch/psviolence00.htm>. Accessed January 17, 2008.
28. Curbow B. Workplace violence: Scope, definition and global context. In: Cooper C and Swanson N, eds. *Workplace Violence in the Health Sector. State of the Art*. Geneva: International Labour Office, International Council of Nurses, World Health Organization and Public Services International, 2002: 35-48.
29. Arnetz JE, Arnetz BB, Petterson I. Violence in the nursing profession: Occupational and lifestyle risk factors in Swedish nurses. *Work & Stress* 1996; 10(2): 119-27.
30. Camerino D, Estryng-Behar M, Conway PM, et al. Work-related factors and violence among nursing staff in the European NEXT study: A longitudinal cohort study. *International Journal of Nursing Studies* 2007; 45(1): 35-50.

31. Duncan SM, Hyndman K, Estabrooks CA, et al. Nurses' experience of violence in Alberta and British Columbia hospitals. *Canadian Journal of Nursing Research* 2001; 32(4): 57-78.
32. Gerberich SG, Church TR, McGovern PM, et al. Risk factors for work-related assaults on nurses. *Epidemiology* 2005; 16(5): 704-9.
33. Leather P. Workplace violence: Scope, definition and global context. In: Cooper C and Swanson N, eds. *Workplace Violence in the Health Sector. State of the Art*. Geneva: International Labour Office, International Council of Nurses, World Health Organization and Public Services International, 2002, 2002: 3-18.
34. Shields M, Wilkins K. *Findings from the 2005 National Survey of the Work and Health of Nurses* (Statistics Canada, Catalogue 83-003-XPE) Ottawa: Minister of Industry, 2006. Available at: <http://www.statcan.ca/english/research/11-621-MIE/11-621-MIE2006052.htm>.
35. Aiken LH, Patrician PA. Measuring organizational traits of hospitals: The Revised Nursing Work Index. *Nursing Research* 2000; 49(3): 146-53.
36. Lake E. Development of the Practice Environment Scale of the Nursing Work Index. *Research in Nursing & Health* 2002; 25: 176-88.
37. Kleim G, Bélanger Y. Using bootstrap variance calculations for a survey with a simple design: The case of the 2005 National Survey of the Work and Health of Nurses. Presented at the Joint Statistical Meetings, Section on Survey Research Methods, August 2007. Salt Lake City, Utah: 2007.
38. Jackson D, Clare J, Mannix J. Who would want to be a nurse? Violence in the workplace—a factor in recruitment and retention. *Journal of Nursing Management* 2002; 10(1): 13-20.
39. Nachreiner NM, Gerberich SG, Ryan AD, et al. Minnesota nurses' study: perceptions of violence and the work environment. *Industrial Health* 2007; 45(5): 672-8.
40. Poster EC, Ryan J. A multiregional study of nurses' beliefs and attitudes about work safety and patient assault. *Hospital & Community Psychiatry* 1994; 45(11): 1104-8.
41. Erickson L, Williams-Evans SA. Attitudes of emergency nurses regarding patient assaults. *Journal of Emergency Nursing* 2000; 26(3): 210-5.
42. Findorff MJ, McGovern PM, Wall MM, et al. Reporting violence to a health care employer: a cross-sectional study. *Journal of the American Association of Occupational Health Nurses* 2005; 53(9): 399-406.
43. Lyneham J. Violence in New South Wales emergency departments. *Australian Journal of Advanced Nursing* 2000; 18(2): 8-17.
44. Rippon TJ. Aggression and violence in health care professions. *Journal of Advanced Nursing* 2000; 31(2): 452-60.
45. Healthcare Commission. Detailed spreadsheets - *National Survey of NHS Staff 2005*. Available at: <http://www.healthcarecommission.org.uk/healthcareproviders/nationalfindings/surveysofnhsstaff/2005.cfm>. Accessed July 17, 2008.
46. Hegney D, Plank A, Parker V. Workplace violence in nursing in Queensland, Australia: a self-reported study. *International Journal of Nursing Practice* 2003; 9(4): 261-8.
47. Whittington R, Shuttleworth S, Hill L. Violence to staff in a general hospital setting. *Journal of Advanced Nursing* 1996; 24(2): 326-33.
48. Whittington R, Wykes T. Violence in psychiatric hospitals: are certain staff prone to being assaulted? *Journal of Advanced Nursing* 1994; 19(2): 219-25.
49. Nolan P, Dallender J, Soares J, et al. Violence in mental health care: the experiences of mental health nurses and psychiatrists. *Journal of Advanced Nursing* 1999; 30(4): 934-41.

Table A
Odds ratios relating workplace climate factors and other selected characteristics to physical assault by a patient over past 12 months, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005

	Adjusted odds ratio	95% confidence interval	
		from	to
Workplace climate factors			
Staffing/Resource adequacy			
First quartile (most adequate) [†]	1.0
Second quartile	1.2	1.0	1.5
Third quartile	1.5*	1.2	1.8
Fourth quartile (least adequate)	2.1*	1.7	2.6
Nurse-physician working relations			
First quartile (most favourable) [†]	1.0
Second quartile	1.1	0.9	1.4
Third quartile	1.1	0.9	1.3
Fourth quartile (least favourable)	1.0	0.8	1.2
Low supervisor support			
Yes	1.2*	1.0	1.4
No [†]	1.0
Low co-worker support			
Yes	1.4*	1.3	1.7
No [†]	1.0
Personal characteristics			
Sex			
Female	0.6*	0.5	0.8
Male [†]	1.0
Years of experience in nursing			
Fewer than 5	1.3*	1.0	1.6
5 to 9	1.1	0.9	1.4
10 to 14	1.1	0.8	1.4
15 to 19 [†]	1.0
20 to 24	0.9	0.7	1.2
25 to 29	0.8	0.7	1.0
30 or more	0.7*	0.6	0.9
Bachelor's degree or higher in nursing			
Yes	1.0	0.8	1.2
No [†]	1.0
General health			
Good, very good or excellent [†]	1.0
Fair or poor	1.0	0.7	1.2
Mental health			
Good, very good or excellent [†]	1.0
Fair or poor	1.4	1.0	1.9
Job satisfaction			
Very satisfied [†]	1.0
Somewhat satisfied	1.1	0.9	1.3
Somewhat dissatisfied	1.5*	1.1	1.9
Very dissatisfied	1.6*	1.0	2.4
Type of nurse			
Registered nurse [†]	1.0
Licensed practical nurse	1.3*	1.1	1.5
Registered psychiatric nurse	1.5*	1.2	1.9
Job characteristics			
Work setting			
Hospital [†]	1.0
Long-term care facility	1.6*	1.3	2.0
Clinical area of employment			
Medicine/Surgery	1.0	0.8	1.3
Psychiatry/Mental health	1.6*	1.2	2.2
Paediatrics	0.5*	0.3	0.9
Maternity/Newborn	0.1*	0.1	0.2
Geriatrics/Long-term care	1.6*	1.2	2.1
Critical care	1.5*	1.1	2.1
Ambulatory care	0.5*	0.3	0.8
Operating/Recovery room	0.5*	0.3	0.7
Emergency room	1.3	1.0	1.7
Several clinical areas [†]	1.0
Oncology	0.5*	0.3	0.9
Rehabilitation	0.8	0.5	1.2
Palliative care	1.4	1.0	2.0
Other direct care	0.8	0.5	1.2
Work status			
Full-time	1.0	0.9	1.2
Part-time [†]	1.0
Shift usually worked			
Days [†]	1.0
Evenings	1.7*	1.3	2.1
Nights	1.8*	1.4	2.2
Mixed	1.7*	1.4	2.0
Length of shift			
12 hours	1.6*	1.4	1.9
Under 12 hours [†]	1.0

[†] reference category

* significantly different from estimate for reference category ($p < 0.05$)

... not applicable

Source: 2005 National Survey of the Work and Health of Nurses.

Table B
Odds ratios relating workplace climate factors and other selected characteristics to emotional abuse from a patient over past 12 months, nurses providing direct care in hospitals or long-term care facilities, Canada, 2005

	Adjusted odds ratio	95% confidence interval	
		from	to
Workplace climate factors			
Staffing/Resource adequacy			
First quartile (most adequate)	1.0
Second quartile	1.3*	1.1	1.6
Third quartile	1.7*	1.4	2.1
Fourth quartile (least adequate)	2.2*	1.8	2.6
Nurse-physician working relations			
First quartile (most favourable) [†]	1.0
Second quartile	1.1	0.9	1.3
Third quartile	1.3*	1.1	1.5
Fourth quartile (least favourable)	1.5*	1.3	1.8
Low supervisor support			
Yes	1.2*	1.0	1.4
No [†]	1.0
Low co-worker support			
Yes	1.3*	1.2	1.5
No [†]	1.0
Personal characteristics			
Sex			
Female	0.9	0.7	1.2
Male [†]	1.0
Years of experience in nursing			
Fewer than 5	0.8*	0.6	1.0
5 to 9	0.8	0.6	1.1
10 to 14	0.9	0.7	1.2
15 to 19 [†]	1.0
20 to 24	0.8	0.6	1.0
25 to 29	0.7*	0.6	0.9
30 or more	0.7*	0.5	0.9
Bachelor's degree or higher in nursing			
Yes	1.1	0.9	1.3
No [†]	1.0
General health			
Good, very good or excellent [†]	1.0
Fair or poor	1.1	0.8	1.5
Mental health			
Good, very good or excellent [†]	1.0
Fair or poor	1.7*	1.3	2.4
Job satisfaction			
Very satisfied [†]	1.0
Somewhat satisfied	1.1	1.0	1.3
Somewhat dissatisfied	1.2	0.9	1.6
Very dissatisfied	1.5	1.0	2.3
Type of nurse			
Registered nurse [†]	1.0
Licensed practical nurse	1.1	0.9	1.2
Registered psychiatric nurse	2.3*	1.9	2.8
Job characteristics			
Work setting			
Hospital [†]	1.0
Long-term care facility	1.1	0.9	1.3
Clinical area of employment			
Medicine/Surgery	1.1	0.9	1.4
Psychiatry/Mental health	2.9*	1.9	4.2
Paediatrics	0.7	0.4	1.2
Maternity/Newborn	0.3*	0.2	0.4
Geriatrics/Long-term care	1.2	0.9	1.6
Critical care	1.1	0.8	1.6
Ambulatory care	1.1	0.7	1.6
Operating/Recovery room	0.4*	0.3	0.6
Emergency room	2.4*	1.8	3.2
Several clinical areas [†]	1.0
Oncology	0.9	0.6	1.3
Rehabilitation	1.0	0.7	1.5
Palliative care	1.0	0.7	1.4
Other direct care	0.9	0.6	1.4
Job tenure			
Full-time	1.0	0.9	1.1
Part-time [†]	1.0
Shift usually worked			
Days [†]	1.0
Evenings	1.4*	1.2	1.8
Nights	1.3*	1.1	1.6
Mixed	1.5*	1.3	1.8
Length of shift			
12 hours	1.6*	1.4	1.9
Under 12 hours [†]	1.0

[†] reference category

* significantly different from estimate for reference category ($p < 0.05$)

... not applicable

Source: 2005 National Survey of the Work and Health of Nurses.