


## Article

# Validation of cognitive functioning categories in the Canadian Community Health Survey—Healthy Aging

by Leanne Findlay, Julie Bernier, Holly Tuokko,  
Susan Kirkland and Heather Gilmour

December 2010



# Validation of cognitive functioning categories in the Canadian Community Health Survey—Healthy Aging

by Leanne Findlay, Julie Bernier, Holly Tuokko, Susan Kirkland and Heather Gilmour

## Abstract

### Background

The objective of this study was to validate categories of cognitive functioning using data from the 2009 Canadian Community Health Survey (CCHS)—Healthy Aging Cognition Module.

### Data and methods

Four measures of cognitive functioning—immediate and delayed recall (memory), and animal-naming and the Mental Alternation Test (executive functioning)—were coded into five categories for the Canadian household population aged 45 or older. The scores for each measure were standardized to t-scores that controlled for age, sex and education. Respondents were classified into five cognitive functioning categories. Cross-tabulations, stratum-specific likelihood ratios and multinomial logit regression were used to assess associations between levels of cognitive functioning and various health outcomes: self-reported general and mental health status, memory and problem-solving ability, activities of daily living, life satisfaction, loneliness, depression, and chronic conditions.

### Results

Results supported the use of five levels of cognitive functioning for all four outcomes on the CCHS—Healthy Aging sample overall and by age group (45 to 64, 65 or older) and language group (English, French).

### Interpretation

These categories can be used in future work on cognitive functioning based on the CCHS—Healthy Aging.

### Keywords

activities of daily living, cognitive disorders, data collection, memory disorders, mental recall, survey methods

### Authors

Leanne Findlay (1-613-951-4648; Leanne.Findlay@statcan.gc.ca), Julie Bernier (1-613-951-4556; Julie.Bernier@statcan.gc.ca) and Heather Gilmour (1-613-951-2114; Heather.Gilmour@statcan.gc.ca) are with the Health Analysis Division at Statistics Canada, Ottawa, Ontario K1A 0T6. Holly Tuokko is with the University of Victoria. Susan Kirkland is with Dalhousie University.

While cognitive decline is not an inevitable consequence of aging, it is more prevalent at older ages.<sup>1</sup> In 2006, one in seven Canadians (13.7% of the total population) was aged 65 or older.<sup>2</sup> Among these seniors, the percentage aged 80 or older continues to grow, as does the number of centenarians. These trends suggest that a rise in the prevalence of cognitive impairment can be anticipated.

Mild cognitive decline heightens the risk of further deterioration,<sup>3-5</sup> but seniors with relatively low levels of impairment may not be identified in cognition studies, which typically focus on people diagnosed with dementia.<sup>6</sup> Nonetheless, a substantial share of the senior population is affected. According to the 1991 Canadian Study of Health and Aging (CSHA), about 17% of Canadians aged 65 or older had mild impairment, often labelled “cognitive impairment—no dementia or CIND.”<sup>7</sup> Similarly, data from the Health and Retirement Survey indicate that 22% of Americans aged 71 or older had CIND.<sup>6</sup> Consequently, examination of the prevalence of various levels of cognitive well-being is warranted.

The last national survey to include measures of cognitive functioning among seniors was the CSHA.<sup>7</sup> The present analysis uses data from the Cognition Module of the 2009 Canadian Community

Health Survey (CCHS)—Healthy Aging to validate a categorization of levels of cognitive functioning in the Canadian household population aged 45 or older. Five categories of four measures of cognitive functioning are examined for the entire sample, and by age group and language.

## Methods

### Data source

The 2009 Canadian Community Health Survey (CCHS)—Healthy Aging is a population-based, cross-sectional survey. The sampling frame consisted of people aged 45 or older living in private dwellings in the ten provinces. The survey excluded residents of the three territories, some remote regions, institutions, Indian reserves or Crown lands, and military bases (military and civilian), and full-time members of the Canadian Forces. Data collection took

place from December 1, 2008 through November 30, 2009 using Computer-Assisted Personal Interviewing.

The purpose of the Cognition Module of the survey was to examine cognitive functioning (as opposed to cognitive impairment) across the lifespan. The Module was administered in English and French to *non-proxy* respondents who consented to participate. This differed from the main component of the survey, for which proxy responses were accepted if the mental or physical health of selected participants prevented them from completing the interview (2.2% of the sample). Preliminary analyses suggested that respondents interviewed by proxy were more likely than those who answered on their own behalf to have dementia or to have suffered a stroke. Exclusion of these respondents from the Cognition Module means that the data may slightly overestimate cognitive functioning in the household population.

The overall response rate to the Cognition Module was 62.3% ( $N = 25,864$ ), compared with 74.4% ( $N = 30,865$ ) for the entire CCHS—Healthy Aging.

### Cognition Module variables

Previous studies have used clinical and non-clinical means to assess cognitive functioning. The Mini-Mental State Examination (MMSE)<sup>8</sup> and the Modified Mini-Mental State Examination (3MS)<sup>9</sup> are the instruments most commonly employed in clinical settings.<sup>10-12</sup> However, when clinical assessment is not possible (in large, survey-based studies), other measures must be used.

Cognition may be defined in terms of domains, including memory and executive functioning (for example, planning, problem-solving, and anticipation of outcomes).<sup>13</sup> The 2009 CCHS—Healthy Aging Cognition Module includes four cognitive tasks: two relating to memory (immediate and delayed recall) and two relating to executive functioning (animal-naming and the Mental Alternation Test). These tasks are similar to those used in other

population-based surveys,<sup>14</sup> as well as in community-based studies.<sup>15-18</sup>

#### Recall tasks

A modified version of the Rey Auditory Verbal Learning Test (RAVLT) was administered to CCHS—Healthy Aging respondents. The test involves memorizing 15 common unrelated words (for example, drum, curtain, bell) and performing two recall trials: one immediate and one delayed. The delayed recall trial took place five minutes after the immediate recall trial (the other cognitive tasks were performed between the recalls). Survey-administered tests of immediate and delayed recall have been shown to be related to each other in a consistent way, to have similar consistency across racial groups,<sup>19</sup> and to have good construct validity.<sup>20</sup>

#### Animal-naming

To test semantic fluency, respondents were given one minute in which to name as many items as possible from a category, in this case, animals. Different types of the same species were counted (for example, robin and parrot counted for two points), but different varieties of the same type (for example, American robin and European robin) received only one point. The animal-naming test has been widely administered, demonstrated to be appropriate for evaluating different populations, and sensitive to different types of brain abnormalities, and it correlates with other tests of verbal fluency.<sup>21</sup>

#### Mental Alternation Test

The Mental Alternation Test (MAT) assesses processing speed.<sup>17,22</sup> Respondents are asked recite the alphabet, and then to count from 1 to 26. They then have 30 seconds in which to alternate between numbers and letters in the sequence 1-A, 2-B, 3-C, etc. The maximum possible score is 51.

Of those who completed any part of the CCHS—Healthy Aging Cognition Module, 85.9% responded to the immediate recall, 75.5% to the delayed recall, 92.6% to animal-naming, and 90.9% to the Mental Alternation Test.

Existing French versions of the recall and animal-naming instruments were used for interviews conducted in French; the English version of the MAT was translated into French.

The various measures reflect independent markers of cognitive functioning, and may have different associations with health outcomes. For example, memory impairment may be important for the early detection of dementia,<sup>23</sup> and declines in executive functioning, as well as memory, may influence activities of daily living.<sup>24</sup> It is also possible that subgroups respond differently to the various measures of cognitive functioning. For instance, people aged 45 to 64 may not demonstrate the same patterns of cognitive functioning as seniors, and patterns may vary by language group.

### Socio-demographic characteristics

Cognitive functioning is typically evaluated in terms of age, sex and education, factors known to be related to cognitive performance.<sup>6,7,10-12,25-28</sup> For instance, results from the English Longitudinal Study of Aging revealed better cognitive performance among younger people, women and individuals with higher education.<sup>14</sup>

Respondents to the CCHS—Healthy Aging reported their sex, age in years and highest level of education. Ten education levels were specified. The language of the interview (English or French) was recorded by the interviewer; respondents who did not complete the CCHS—Healthy Aging in either English or French were excluded from the Cognition Module.

### Analysis variables

Numerous physical and psychological correlates of impaired cognitive functioning have been identified (Table 1). Cognitive difficulties have been associated with lower self-rated health,<sup>29,30</sup> depression,<sup>31-33</sup> loneliness,<sup>34,35</sup> decreased life satisfaction,<sup>36,37</sup> and reduced ability to perform instrumental activities of daily living.<sup>24,38-40</sup> People with

*Validation of cognitive functioning categories in the Canadian Community Health Survey • Methodological insights*

**Table 1**  
**Selected characteristics of 2009 Canadian Community Health Survey—**  
**Healthy Aging Cognition Module respondents, household population aged 45**  
**or older, Canada excluding territories**

Characteristic	%	Mean	Standard error
<b>Socio-demographic</b>			
<b>Sex</b>			
Women	51.9	...	...
Men	48.1	...	...
<b>Age</b>			
	...	60.5	0.04
<b>Marital status</b>			
Married	73.5	...	...
Single	26.5	...	...
<b>Language</b>			
English	76.7	...	...
French	23.3	...	...
<b>Education</b>			
Grade 8 or lower (Quebec Secondary II or lower)	9.2	...	...
Grades 9 or 10	8.0	...	...
Grades 11 to 13	3.9	...	...
Secondary graduation	19.1	...	...
Some postsecondary	5.4	...	...
Trades certificate/diploma	13.7	...	...
College/CEGEP certificate/diploma	17.1	...	...
University certificate below bachelor's degree	3.2	...	...
Bachelor's degree	12.2	...	...
University certificate above bachelor's degree	8.3	...	...
<b>Province</b>			
Newfoundland and Labrador	1.7	...	...
Prince Edward Island	0.4	...	...
Nova Scotia	3.1	...	...
New Brunswick	2.4	...	...
Quebec	24.6	...	...
Ontario	38.3	...	...
Manitoba	3.4	...	...
Saskatchewan	2.9	...	...
Alberta	9.3	...	...
British Columbia	13.8	...	...
<b>Cognitive outcomes</b>			
Immediate recall	...	5.5	0.10
Delayed recall	...	4.0	0.12
Animal naming	...	17.9	0.03
Mental Alternation Test	...	22.6	0.03
<b>Health outcomes</b>			
<b>Self-perceived health</b>			
Excellent/Very good/Good	84.8	...	...
Fair/Poor	15.2	...	...
<b>Self-perceived mental health</b>			
Excellent/Very good/Good	94.2	...	...
Fair/Poor	5.8	...	...
<b>Life satisfaction</b>			
Not low	84.4	...	...
Low	15.6	...	...
<b>Likelihood of depression</b>			
Less than 0.9 probability	94.3	...	...
0.9 probability or higher	5.7	...	...
<b>Loneliness</b>			
Not high	87.8	...	...
High	12.2	...	...
<b>Activities of daily living</b>			
No problems	90.2	...	...
Mild/Moderate/Severe/Total problems	9.8	...	...
<b>Memory</b>			
Able to remember most things	75.9	...	...
Somewhat forgetful/Very forgetful/Unable to remember anything	24.1	...	...
<b>Ability to think clearly and solve problems</b>			
Able to think clearly/solve problems	91.9	...	...
Having a little/some/great deal of difficulty	8.1	...	...
<b>Chronic conditions</b>			
Neurological disorder	2.7	...	...
Vascular disorder	44.1	...	...
Psychiatric disorder	10.5	...	...

... not applicable

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

cerebrovascular disease,<sup>29</sup> diabetes,<sup>33,41-44</sup> hypertension,<sup>45</sup> or stroke<sup>12,46</sup> are more likely to be cognitively impaired than are individuals without these conditions. Psychiatric disorders have also been associated with poor cognitive functioning.<sup>47,48</sup>

*Self-perceived health*

CCHS—Healthy Aging respondents were asked about their general and mental health: “In general, would you say your [mental] health is. . . .” The response options—“excellent,” “very good,” “good,” “fair,” and “poor”—were dichotomized to reflect good (excellent/very good/good) versus poor (fair/poor) health.

*Activities of daily living*

Questions about respondents’ ability to perform activities of daily living (ADL) were based on the OARS Multidimensional Assessment Questionnaire.<sup>49</sup> An overall summary measure of ratings on the ADL capacity-instrumental and physical dimensions was derived. A score of 0 indicates no functional impairment; 1 = mild impairment; 2 = moderate impairment; 3 = severe impairment; and 4 = total impairment. Responses were dichotomized to identify respondents with no impairment versus mild/moderate/severe/total impairment.

*Life satisfaction*

On a scale from 0 to 10, with 0 representing very dissatisfied and 10, very satisfied, respondents were asked: “How do you feel about your life as a whole right now?” Scores were dichotomized to identify those whose life satisfaction was low (at least 1 standard deviation below the mean) versus not low.

*Depression*

The CCHS—Healthy Aging measure of depression is a shortened version of the World Health Organization Composite International Diagnostic Interview (CIDI) Scale, which is based on the DSM-III-R and the Diagnostic Criteria for the Research of the ICD-10. The depression subscale pertains to people who felt depressed or lost interest in

things for two or more weeks in the past year. For the CCHS—Healthy Aging, a derived variable was created based on the depression score, indicating the probability that respondents would have been diagnosed as having experienced a major depressive episode in the past 12 months if they had completed the Long-Form CIDI. A probability of 0 was assigned to respondents who replied negatively to the stem question (did not have depression for two or more weeks in the past year); a cut-off value of 0.9 was used to distinguish those with a high probability of depression (above 0.9) from those with a lower probability.

### *Loneliness*

The 3-Item Loneliness Scale<sup>50</sup> measures an individual's reported loneliness. On a 3-point Likert scale (“hardly ever,” “some of the time,” and “often”), CCHS—Healthy Aging respondents answered the questions: “How often do you feel:

- that you lack of companionship?”
- left out?”
- isolated from others?”

Higher scores indicate greater loneliness. Scores were dichotomized to identify those with high loneliness (at least 1 standard deviation above the mean) versus not high loneliness.

### *Self-reported cognition (Health Utilities Index)*

The Health Utilities Index (HUI) Mark III assesses functional health status in eight domains: vision, hearing, speech, ambulation, dexterity, emotion, cognition, and pain.<sup>51,52</sup> The HUI has been shown to have strong reliability and validity in general,<sup>53</sup> as well as for patients with lower cognitive functioning.<sup>54</sup>

Only the cognition subscale of the HUI was pertinent to the current study. The items of interest were: “How would you describe your usual ability to:

- remember things (able to remember most things; somewhat forgetful; very forgetful; unable to remember anything at all).”
- think and solve day-to-day problems (able to think clearly and solve

problems; having a little difficulty; having some difficulty; having a great deal of difficulty; unable to think or solve problems).”

Items were dichotomized as “able to remember most things” versus at least “somewhat forgetful,” and “able to think clearly and solve problems” versus having “at least some difficulty.”

### *Chronic conditions*

Respondents were asked if they had been diagnosed with specific long-term health conditions. Conditions relevant to the current analysis were grouped into neurological (Alzheimer's Disease or other dementia, Parkinson's Disease, effects of stroke), vascular (high blood pressure, diabetes, heart attack, heart disease), and psychiatric (mood disorder or anxiety) disorders.

### **Analytical techniques**

#### *T-score creation*

Selecting cut-points to identify impairment implies that definitive lines demarcate “normal” from “dysfunctional” scores. It is more likely that cognitive functioning operates on a continuum and that several categories are more appropriate as indicators of impairment.<sup>4</sup> Consequently, for this analysis, multiple categories of cognitive functioning were identified.<sup>55</sup> T-scores that control for age, sex and education can be calculated for each cognitive measure. Using the sample data for the current study, five categories of cognitive functioning were created, representing t-scores of 0 to 34, 35 to 44, 45 to 54, 55 to 64, and 65 or more.

To generate t-scores from the results of each of the four cognitive tasks, raw scores were converted to scaled scores (mean = 10, standard deviation = 3); higher scaled scores indicate better performance.<sup>55</sup> Scaled scores were regressed separately for each task on age, sex and education. In this manner, equations were created for each dependent variable (cognitive outcome) in the form:

$$DV = \text{intercept} + b(\text{age}) + b(\text{education}) + b(\text{gender})$$

Each respondent's predicted scaled score was generated from this equation (that is, independent of age, sex and education). The respondent's predicted score was subtracted from the actual scaled score to determine the residual, indicating how well the individual performed, compared with what would be expected based on his/her age, sex and education. Finally, residual scores were converted to t-scores with the following equation:

$$T = \{[(\text{residual}/\text{standard deviation of the residual}) \times 10] + 50\}$$

Thus, the t-scores are independent of age, sex and education; are normally distributed; have a mean of 50 and a standard deviation of 10; and are independent of a unit of measurement.<sup>55</sup>

#### *Validation*

Once t-scores were created and individuals were assigned to one of the five cognitive function categories, the first step in empirically validating the categories was to examine cross-tabulations of the categories by health outcome.

Stratum-specific likelihood ratios (SSLRs) were calculated to determine the accuracy of assigning individuals to levels of cognitive functioning based on the health outcomes.<sup>56-58</sup> SSLRs are generalizable and independent of actual probabilities in the population.<sup>59</sup> The likelihood that people in each cognitive functioning category (stratum) will experience a certain outcome (for example, fair/poor self-rated health) is given relative to their likelihood of experiencing a positive outcome (in this example, excellent/very good/good self-rated health), according to the formula:

$$SSLR = (x_{1g}/n_1)/(x_{0g}/n_0)$$

where  $x_{1g}$  is the number of people with the health outcome (fair/poor health) in the  $g$ th stratum;  $n_1$  is the total number of people with the health outcome;  $x_{0g}$  is the number of people without the health outcome (in good health) in the

**Validation of cognitive functioning categories in the Canadian Community Health Survey • Methodological insights**

gth stratum; and  $n_0$  is the total number of people without the health outcome.

An SSLR of 10 or more indicates that the health outcome is highly likely; an SSLR below 0.1 indicates that it is highly unlikely.<sup>56,57</sup> It is anticipated that SSLRs would be high when a poor health outcome is more likely, and low when a poor health outcome is unlikely.

The final step was to examine all relevant health variables as predictors of cognitive functioning, comparing lower levels of functioning to the highest category (t-scores of 65 or more) for each cognitive outcome. Because the dependent variable (cognitive functioning category) comprised five levels, a multinomial logit regression analysis was used. The odds of reporting a health problem (for example, fair/poor health) should be greatest for those in the lowest (versus the highest) cognitive functioning category, with odds decreasing for those in progressively higher categories of functioning.

Results are presented only for the immediate recall outcome; results were similar for delayed recall, animal-naming, and the MAT (Appendix Tables A to I). Correlations between the four outcome variables were moderate (immediate recall with delayed recall,  $r = .69$ ; immediate recall with animal-naming,  $r = .36$ , immediate recall with MAT,  $r = .34$ ; delayed recall with animal-naming,  $r = .33$ , delayed recall with MAT,  $r = .30$ ; animal-naming with MAT,  $r = .45$ ; all  $p$ 's  $\leq .001$ ). Survey sampling weights were applied to all point-estimates to account for the complex survey design of the CCHS—Healthy Aging.

Because the response rate for the Cognition Module was lower than that for the full CCHS—Healthy Aging, separate sampling weights were created for use with the cognitive outcome variables sample. All analyses were performed with SAS 9.1 and SAS-callable SUDAAN. Standard errors in modelling were computed using a bootstrapping technique.<sup>60</sup>

**Results**

**Descriptive statistics**

As expected,<sup>55</sup> the distribution of immediate recall scores across cognitive functioning categories was normal, with the most common category (39% of respondents) being t-scores in the 45-to-54 range (Table 2). Approximately 6% of respondents scored in the lowest category, and 8%, in the highest.

People who reported fair/poor general health were more likely to have relatively low immediate recall scores. About 9% of them had scores in the lowest category, compared with 6% overall. Conversely, 5% of those with fair/poor health had immediate recall scores in the in the highest category, compared with 8% overall. This pattern was even more pronounced for self-reported mental health. Similarly, relatively high

**Table 2**  
**Percentage distribution of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by immediate recall score and selected health characteristics, household population aged 45 or older, Canada excluding territories**

Health characteristic	Immediate recall t-score				
	Low 0 to 34	35 to 44	45 to 54	55 to 64	High 65 or more
	%				
<b>Total</b>	5.6	24.5	39.2	23.0	7.6
<b>Self-perceived health</b>					
Fair/Poor	8.5	28.6	39.8	18.1	5.1
Excellent/Very good/Good	5.1	23.8	39.1	23.8	8.1
<b>Self-perceived mental health</b>					
Fair/Poor	14.9 <sup>F</sup>	32.2	32.3	16.6	3.9 <sup>F</sup>
Excellent/Very good/Good	5.1	24.1	39.6	23.4	7.9
<b>Activities of daily living</b>					
No problems	5.3	24.1	39.2	23.5	7.9
Mild/Moderate/Severe/Total problems	8.6	28.4	39.3	18.4	5.3 <sup>F</sup>
<b>Life satisfaction</b>					
Low	7.8	30.7	38.2	18.0	5.3
Not low	5.2	23.4	39.4	23.9	8.0
<b>Depression</b>					
0.9 probability or higher	5.8 <sup>E</sup>	25.7	39.3	18.3	10.8 <sup>E</sup>
Less than 0.9 probability	5.5	24.4	39.3	23.4	7.4
<b>Loneliness</b>					
High	9.1	28.4	36.8	19.5	6.3
Not high	5.1	24.0	39.6	23.5	7.8
<b>Memory</b>					
Able to remember most things	5.0	24.0	39.5	23.6	7.9
Somewhat forgetful/Very forgetful/Unable to remember anything	7.6	26.4	38.2	21.0	6.8
<b>Ability to think clearly and solve problems</b>					
Able to think clearly/solve problems	5.2	23.8	39.4	23.7	7.9
Having a little/some/great deal of difficulty/unable	10.5	33.9	36.6	14.8	4.3 <sup>F</sup>
<b>Neurological disorder</b>					
Yes	9.0 <sup>F</sup>	30.3	41.1	16.0	F
No	5.6	24.4	39.1	23.2	7.7
<b>Vascular disorder</b>					
Yes	5.0	24.0	41.0	23.0	7.0
No	6.1	25.0	37.9	22.9	8.1
<b>Psychiatric disorder</b>					
Yes	9.6 <sup>E</sup>	26.3	37.6	19.3	7.2 <sup>E</sup>
No	5.2	24.3	39.4	23.4	7.7

<sup>E</sup> Interpret with caution (coefficient of variation 16.6% to 33.3%)

<sup>F</sup> Too unreliable to be reported (coefficient of variation greater than 33.3%)

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

percentages of people who reported difficulties with activities of daily living, lower life satisfaction and loneliness had low immediate recall scores. By contrast, no pattern emerged for depression.

The HUI cognitive functioning variables were associated with immediate recall scores. Respondents who reported that they were at least somewhat forgetful and who had at least some difficulties thinking clearly and solving problems were more likely than others to have immediate recall scores in the lowest category and less likely to have scores in the highest category.

The presence of a neurological or psychiatric disorder was related to cognitive functioning. Relatively high percentages of people who reported such conditions had low immediate recall scores. However, this was not the case for people with vascular disorders.

### Stratum-specific likelihood ratios

In general, the stratum-specific likelihood ratios (SSLR) supported the cognitive functioning categories: the higher their immediate recall score, the less likely were respondents to have negative health characteristics. (Although the SSLR patterns were generally as anticipated,

some differences emerged for delayed recall, animal-naming and MAT—Appendix Tables D to F).

SSLRs for fair/poor self-rated general and mental health, difficulties with activities of daily living, low life satisfaction and loneliness decreased as immediate recall scores rose (Table 3). In general, depression also followed the expected pattern, with the highest SSLR for the lowest immediate recall score category. The two HUI cognition variables also demonstrated the anticipated pattern.

Similarly, the SSLRs for neurological and psychiatric disorders followed the expected pattern in that the likelihood of the conditions was associated with low immediate recall scores; no association was shown for vascular disorders.

### Multinomial logistic regression

The final step was to examine the odds of being in a low immediate recall score category given the presence of a negative health characteristic. The highest t-score category was set as the reference group (Table 4). As expected, scoring in the lowest immediate recall category was associated with the highest odds of poor health. For instance, compared with

people whose immediate recall scores were in the highest category, those with scores in the lowest category had more than twice the odds of being in fair/poor general health, almost six times the odds of being in fair/poor mental health, and more than twice the odds of having difficulties with activities of daily living. Results were similar for low life satisfaction and loneliness. Not surprisingly, people with the lowest immediate recall scores had almost twice the odds of reporting that they were at least somewhat forgetful, and almost four times the odds of reporting that they had some difficulty thinking clearly and solving problems, compared with people with the highest immediate recall scores. However, no association was shown between depression and immediate recall scores.

People with immediate recall scores in the lowest category had more than three times the odds of reporting a neurological condition and twice the odds of reporting a psychiatric disorder. (Odds ratios were generally similar for the other three measures of cognitive functioning, with the exception of psychiatric disorders and the MAT—Appendix Tables G, H and I).

**Table 3**  
**Stratum-specific likelihood ratios (SSLR) for selected health characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by immediate recall score, household population aged 45 or older, Canada excluding territories**

Health characteristic	Immediate recall t-score														
	0 to 34			35 to 44			45 to 54			55 to 64			65 or more		
	SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval	
	from	to	from	to	from	to	from	to	from	to	from	to	from	to	
Low self-rated health	1.66	1.65	1.67	1.20	1.19	1.20	1.02	1.01	1.02	0.76	0.76	0.76	0.63	0.62	0.63
Low self-rated mental health	2.93	2.92	2.95	1.34	1.34	1.34	0.82	0.81	0.82	0.71	0.71	0.72	0.50	0.49	0.51
Difficulties with activities of daily living	1.62	1.61	1.63	1.18	1.17	1.18	1.00	1.00	1.00	0.79	0.78	0.79	0.67	0.66	0.67
Low life satisfaction	1.49	1.49	1.50	1.31	1.31	1.31	0.97	0.97	0.97	0.75	0.75	0.75	0.66	0.66	0.66
High probability of depression	1.07	1.06	1.08	1.05	1.05	1.06	1.00	1.00	1.00	0.78	0.78	0.79	1.46	1.45	1.47
High loneliness	1.79	1.78	1.80	1.18	1.18	1.19	0.93	0.93	0.93	0.83	0.82	0.83	0.80	0.80	0.81
Unable to remember things	1.51	1.50	1.52	1.10	1.10	1.10	0.97	0.97	0.97	0.89	0.89	0.89	0.86	0.86	0.87
Unable to think/solve problems	2.00	1.99	2.01	1.43	1.42	1.43	0.93	0.93	0.93	0.62	0.62	0.63	0.54	0.53	0.54
Neurological disorder	1.61	1.59	1.63	1.24	1.24	1.25	1.05	1.05	1.05	0.69	0.68	0.69	0.47	0.47	0.48
Vascular disorder	0.82	0.82	0.82	0.96	0.96	0.96	1.08	1.08	1.08	1.00	1.00	1.01	0.86	0.86	0.86
Psychiatric disorder	1.86	1.85	1.87	1.08	1.08	1.09	0.95	0.95	0.96	0.82	0.82	0.83	0.94	0.93	0.94

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

**Table 4**  
**Odds ratios relating selected health status characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by immediate recall score, household population aged 45 or older, Canada excluding territories**

Health characteristic	Adjusted degrees of freedom	Adjusted chi-square	Immediate recall t-score											
			0 to 34			35 to 44			45 to 54			55 to 64		
			Odds ratio	95% confidence interval		Odds ratio	95% confidence interval		Odds ratio	95% confidence interval		Odds ratio	95% confidence interval	
from	to	from		to	from		to	from		to				
Low self-rated health	3.59	32.89	2.64	1.64	4.26	1.90	1.35	2.68	1.62	1.15	2.26	1.21	0.85	1.71
Low self-rated mental health	3.17	45.51	5.86	3.10	11.08	2.68	1.72	4.17	1.63	1.04	2.56	1.42	0.86	2.35
Difficulties with activities of daily living	3.63	25.65	2.42	1.51	3.90	1.76	1.19	2.59	1.50	1.03	2.18	1.17	0.79	1.74
Low life satisfaction	3.62	35.90	2.27	1.45	3.53	1.99	1.45	2.72	1.47	1.10	1.97	1.14	0.84	1.54
High probability of depression	3.71	6.19	0.73	0.39	1.39	0.72	0.43	1.23	0.69	0.41	1.15	0.54	0.31	0.92
High loneliness	3.93	27.73	2.23	1.47	3.39	1.48	1.05	2.06	1.16	0.83	1.61	1.03	0.74	1.44
Unable to remember things	3.84	16.23	1.76	1.20	2.58	1.28	0.97	1.69	1.12	0.86	1.47	1.03	0.77	1.38
Unable to think/solve problems	3.71	45.13	3.71	1.94	7.08	2.64	1.45	4.83	1.72	0.95	3.13	1.16	0.62	2.15
Neurological disorder	3.87	17.69	3.40	1.47	7.86	2.62	1.20	5.74	2.22	1.01	4.84	1.45	0.66	3.21
Vascular disorder	3.83	8.94	0.95	0.72	1.26	1.12	0.92	1.36	1.26	1.05	1.51	1.17	0.95	1.43
Psychiatric disorder	3.73	16.52	1.99	1.17	3.36	1.16	0.78	1.73	1.02	0.69	1.50	0.88	0.58	1.33

Note: Comparison group is 65 or more t-score category.

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

### Subgroup analyses

Validation conducted for people aged 45 to 64 and for those aged 65 or older, as well as for English and French respondents, yielded results similar to those obtained for the entire sample (results available upon request). Whether they were in the younger or older age group, English or French, respondents demonstrated similar patterns between health outcomes and the five categories of cognitive functioning (in both cross-tabulations and SSLR comparisons).

Regardless of their age group, people with low immediate recall scores were more likely to have fair/poor self-rated general and mental health, difficulties with activities of daily living, low life satisfaction, loneliness, less ability to think and solve problems, and neurological disorders, compared with people whose scores placed them in higher immediate recall categories. The only differences between the younger and older cohort were in memory and psychiatric disorders—lower immediate recall scores were not strongly associated with ability to remember things and psychiatric disorders among 45- to 64-year-olds, but they were for seniors.

For English and French respondents, lower immediate recall scores were associated with fair/poor self-rated general and mental health, difficulties with activities of daily living, low life satisfaction, loneliness, lower self-rated cognition, neurological disorders, and psychiatric disorders. Depression and vascular disorders were not associated with immediate recall scores for either language group

### Discussion

The results of the current study confirm that categories of cognitive functioning can be described from the CCHS—Healthy Aging Cognition Module. Four tests of cognitive functioning—immediate recall, delayed recall, animal-naming and the Mental Alternation Test—were validated based on literature-supported correlates of cognitive functioning. Lower cognitive functioning (notably, t-scores less than 34) was associated with poorer self-rated general and mental health, difficulties with activities of daily living, lower life satisfaction, and loneliness. As might be expected, self-reported cognitive difficulties (forgetfulness and difficulty

thinking clearly and solving problems) were associated with low immediate recall scores. The fact that the strongest correlates of the cognitive functioning categories were self-rated mental health and difficulties thinking clearly and solving problems lends the greatest support to the use of the categories presented in this analysis.

Cognitive functioning was not associated with the probability of depression. However, the literature on this subject is inconsistent. Some studies have found no association between depression and cognition,<sup>61</sup> while others have shown a relationship, even accounting for socio-economic factors.<sup>33</sup> Beirman et al.<sup>31</sup> suggested a non-linear relationship between depression and cognitive decline, with elevated levels of depression (and anxiety) in the early stages of decline, but diminished levels as deterioration progresses. Further research on the association between depression and cognitive functioning is obviously necessary.

While neurological and psychiatric disorders were associated with lower cognitive functioning, no patterns emerged for vascular disorders. Previous work, too, has suggested that heart



disease, hypertension and diabetes are not necessarily associated with cognitive decline, especially over and above other risk factors such as low educational attainment.<sup>33,62,63</sup>

### **Limitations and future directions**

A major strength of the current study is the large, nationally representative sample. However, several limitations should be acknowledged.

Proxy responses were not accepted for the Cognition Module. Other research has shown that individuals for whom proxy responses are provided tended to perform poorly on cognitive measures and were more likely to have dementia.<sup>20</sup> Thus, the CCHS—Healthy Aging data may underestimate the prevalence of lower cognitive functioning in the Canadian household population.

The CCHS—Healthy Aging Cognition Module used non-clinical measures of cognitive functioning. A clinical assessment would have allowed a test of sensitivity and specificity of the measures in identifying cognitive decline or dementia. This may explain why relationships were not found between vascular disorders (and/or depression) and the cognition categories. Muller et al.<sup>64</sup> found a significant relation between cardiovascular disease and MMSE scores, but not administered tests.

The longitudinal assessment of cognitive functioning among the population is warranted. Such studies would allow researchers to focus on associations between specific risk factors (or correlates) and cognitive functioning over time. For instance, Wilson et al.<sup>36</sup> found that loneliness was associated with a more rapid cognitive decline in elderly people.

### **Conclusions**

Based on the results of tests of immediate and delayed recall, animal-naming, and the MAT in the CCHS—Healthy Aging Cognition Module, five categories describing low to high cognitive functioning were created. These categories were validated for the household population aged 45 or older overall, and by age group and language.

The aging of Canada's population will likely be accompanied by a growing number of people experiencing cognitive decline. CCHS—Healthy Aging data can contribute an understanding of the prevalence of this condition in the

household population. This validation study enhances the analytic value of the information in the Cognition Module. ■

### **Acknowledgements**

Statistics Canada thanks all participants for their input and advice during the development of the Canadian Community Health Survey—Healthy Aging. The survey content was developed by the Health Statistics Division at Statistics Canada in consultation with Health Canada, the Public Health Agency of Canada, and experts conducting the Canadian Longitudinal Study on Aging (CLSA), a major strategic initiative of the Canadian Institute of Health Research. Consultations included stakeholders from Human Resources and Social Development Canada and provincial and territorial health ministries. The addition of 5,000 respondents aged 45 to 54 was funded by the CLSA.

# References

1. Rowe JW, Khan RL. Human aging: usual and successful. *Science* 1987; 237: 143-9.
2. Statistics Canada. *Portrait of the Canadian Population in 2006, by Age and Sex, 2006 Census* (Catalogue 97-551-XIE) Ottawa: Minister of Industry, 2007.
3. Matthews FE, Stephan BCM, McKeith IG, et al. Two-year progression from mild cognitive impairment to dementia: To what extent do different definitions agree? *Journal of the American Geriatric Society* 2008; 56: 1424-33.
4. Tuokko H, Frerichs RJ. Cognitive Impairment with No Dementia (CIND): Longitudinal studies, the findings, and the issues. *The Clinical Neuropsychologist* 2000; 14: 504-25.
5. Tuokko H, Hultsch DF. *Mild Cognitive Impairment: International Perspectives*. New York: Taylor and Francis, 2006.
6. Plassman BL, Langa KM, Fisher GG, et al. Prevalence of cognitive impairment without dementia in the United States. *Annals of Internal Medicine* 2008; 148(427): 434.
7. Graham JE, Rockwood K, Beattie BL, et al. Prevalence and severity of cognitive impairment with and without dementia in an elderly population. *Lancet* 1997; 349: 1793-6.
8. Folstein MF, Folstein SE, McHugh PR. 'Mini-Mental State': a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research* 1975; 12: 189-98.
9. Teng EL, Chui HC. The Modified Mini-Mental State (3MS) examination. *Journal of Clinical Psychiatry* 1987; 48: 314-8.
10. Bravo G, Hébert R. Age- and education-specific reference values for the Mini-Mental and Modified Mini-Mental State Examinations derived from a non-demented elderly population. *International Journal of Geriatric Psychiatry* 1997; 12: 1008-18.
11. Crum R, Anthony JC, Bassett SS, et al. Population-based norms for the Mini-Mental State Examination by age and educational level. *Journal of the American Medical Association* 1993; 269(18): 2386-91.
12. McDowell I, Xi G, Lindsay J, et al. Canadian Study of Health and Aging: Study description and patterns of early cognitive decline. *Aging Neuropsychology and Cognition* 2004; 11(2-3): 149-68.
13. Lezak MD, Howieson DB, Loring DW, et al. *Neuropsychological Assessment: 4th Edition*. New York: Oxford University Press, Inc., 2004.
14. Marmot M, Banks J, Blundell R, et al. *Health, Wealth and Lifestyles of the Older Population in England: The 2002 English Longitudinal Study of Ageing*. London, UK: Institute for Fiscal Studies, 2003.
15. Baars MAE, Boxtel MPJ, Dijkstra JP, et al. Predictive value of mild cognitive impairment for dementia. *Dementia and Geriatric Cognitive Disorders* 2009; 27: 173-81.
16. Carlson MC, Xue Q, Zhou J, et al. Executive decline and dysfunction precedes declines in memory: The women's health and aging study 2. *Journal of Gerontology: Medical Sciences* 2009; 64: 110-7.
17. Salib E, McCarthy J. Mental alteration test (MAT): A rapid and valid screening tool for dementia in primary care. *International Journal of Geriatric Psychiatry* 2002; 17: 1157-61.
18. Billick SB, Siedenbueg E, Burgett W, et al. Validation of the Mental Alteration Test with the Mini-Mental State Examination in geriatric psychiatric inpatients and normal controls. *Comprehensive Psychiatry* 2001; 42(3): 202-5.
19. Ofstedal MB, Fisher GG, Herzog AR. *HRS/AHEAD Documentation Report*. Ann Arbor, MI: University of Michigan, 2005.
20. Herzog AR, Wallace RB. Measures of cognitive functioning in the AHEAD Study. *Journals of Gerontology Series B, Psychological Sciences and Social Sciences* 1997; 52 (special issue): 37-48.
21. Ardilia A, Ostrosky-Solis F, Bernal B. Cognitive testing toward the future: The example of Semantic Verbal Fluency (ANIMALS). *International Journal of Psychology* 2006; 41(5): 324-32.
22. Teng EL. The Mental Alteration Test (MAT). *The Clinical Neuropsychologist* 1995; 9: 287.
23. Tuokko H, Vernon-Wilkinson R, Weir J, et al. Cued recall and early identification of dementia. *Journal of Clinical and Experimental Neuropsychology* 1991; 13: 871-9.
24. Tomaszewski Farias S, Cahn-Weiner DA, Harvey DJ, et al. Longitudinal changes in memory and executive functioning are associated with longitudinal change in instrumental activities of daily living in older adults. *The Clinical Neuropsychologist* 2009; 23: 446-61.
25. Ratcliffe G, Dodge H, Birzescu M, et al. Tracking cognitive functioning over time: Ten-year longitudinal data from a community-based sample. *Applied Neuropsychology* 2003; 10(2): 76-88.
26. Tombaugh T, Kozak J, Rees L. Normative data stratified by age and education for two measures of verbal fluency: FAS and Animal Naming. *Archives of Clinical Neuropsychology* 1999; 14(2): 167-77.
27. Jones TG, Schinka JA, Vanderplaeg RD, et al. 3MS normative data for the elderly. *Archives of Clinical Neuropsychology* 2002; 17: 171-7.
28. Tucker-Drob EM, Johnson KE, Jones RN. The cognitive reserve hypothesis: A longitudinal examination of age-associated declines in reasoning and processing speed. *Developmental Psychology* 2009; 45: 431-46.
29. Frisoni GB, Fratiglioni L, Fastbom J, et al. Mild cognitive impairment in the population and physical health: Data on 1,435 individuals aged 75 to 95. *The Journals of Gerontology* 2000; 55A(6): M322-8.
30. Walker JD, Maxwell CJ, Hogan DB, et al. Does self-rated health predict survival in older persons with cognitive impairment? *Journal of the American Geriatric Society* 2004; 52: 1895-900.
31. Bierman EJM, Comijs HC, Jonker C, et al. Symptoms of anxiety and depression in the course of cognitive decline. *Dementia and Geriatric Cognitive Disorders* 2007; 24: 213-9.
32. Bruce JM, Bhalla R, Westervelt HJ, et al. Neuropsychological correlates of self-reported depression and self-reported cognition among patients with mild cognitive impairment. *Journal of Geriatric Psychiatry and Neurology* 2008; 21(1): 34-40.
33. Lopez OL, Jagust WJ, Dulberg C, et al. Risk factors for mild cognitive impairment in the cardiovascular health study cognition study. *Archives of Neurology* 2003; 60: 1394-9.
34. Holmen K, Ericsson K, Winblad B. Social and emotional loneliness among non-demented and demented elderly people. *Archives of Gerontology and Geriatrics* 2000; 31: 177-92.
35. Wilson RS, Krueger KR, Arnold SE, et al. Loneliness and risk of Alzheimer disease. *Archives of General Psychiatry* 2007; 64: 234-40.
36. Jones TG, Rapport LJ, Hanks RA, et al. Cognitive and psychosocial predictors of subjective well-being in urban older adults. *Clinical Neuropsychologist* 2003; 17: 3-18.
37. Berg AI, Hassing LB, McClearn GE, et al. What matters for life satisfaction in the oldest-old? *Aging and Mental Health* 2006; 10: 257-64.
38. Nourhashemi F, Andrieu S, Gillette-Guyonnet S, et al. Instrumental activities of daily living as a potential marker of frailty: A study of 7364 community-dwelling elderly women (the EPIDOS Study). *The Journals of Gerontology* 2001; 56A: M448-53.
39. Pereira FS, Yassuda MS, Oliveira AM, et al. Executive dysfunction correlates with impaired functional status in older adults with varying degrees of cognitive impairment. *International Psychogeriatrics* 2008; 20: 1104-15.

40. Razani J, Casas R, Wong J, et al. The relationship between executive functioning and activities of daily living in patients with relatively mild dementia. *Applied Neuropsychology* 2007; 14: 1-7.
41. Beissels GJ, Staekenborg S, Brunner E, et al. Risk of dementia in diabetes mellitus: a systematic review. *Lancet Neurology* 2006; 5: 64-74.
42. Strachan MWJ, Ewing FME, Deary IJ, et al. Is Type II diabetes associated with an increased risk in cognitive dysfunction? *Diabetes Care* 1997; 20(3): 438-45.
43. van den Berg E, de Craen AJM, Biessel GJ, et al. The impact of diabetes mellitus on cognitive decline in the oldest of the old: a prospective population-based study. *Diabetologia* 2006; 49: 2015-23.
44. Luck T, Riedel-Heller SG, Kaduszkiewicz H, et al. Mild cognitive impairment in general practice: Age-specific prevalence and correlate results from the German Study on Ageing, Cognition and Dementia in Primary Care Patients (AgeCoDe). *Dementia and Geriatric Cognitive Disorders* 2007; 24: 307-16.
45. Cherubini A, Lowenthal DT, Paran E, et al. Hypertension and cognitive function in the elderly. *Dis Mon* 2010; 56: 106-47.
46. Rafnsson SB, Deary IJ, Smith FB, et al. Cardiovascular diseases and decline in cognitive function in an elderly community population: The Edinburgh artery study. *Psychosomatic Medicine* 2007; 69: 425-34.
47. Colenda CC, Legault C, Rapp SR, et al. Psychiatric disorders and cognitive dysfunction among older, postmenopausal women: Results from the women's health initiative memory study. *American Journal of Geriatric Psychiatry* 2010; 18: 177-86.
48. Rosenberg PB, Mielke MM, Xue Q, et al. Depressive symptoms predict incident cognitive impairment in cognitively healthy older women. *American Journal of Geriatric Psychiatry* 2010; 18: 204-11.
49. Fillenbaum GG, Smyer MA. The development, validity, and reliability of the Oars Multidimensional Functional Assessment Questionnaire. *Journal of Gerontology* 1981; 36: 428-34.
50. Hughes ME, Waite LJ, Hawkey LC, et al. A short scale for measuring loneliness in large surveys: Results from two population based studies. *Research on Aging* 2004; 26: 655-72.
51. Feeney D, Furlong W, Torrance GW, et al. Multiattribute and single-attribute utility functions for the Health Utilities Index mark 3 system. *Medical Care* 2002; 40: 113-28.
52. Feng Y, Bernier J, McIntosh C, et al. Validation of disability categories derived from Health Utilities Index Mark 3 scores. *Health Reports (Statistics Canada, Catalogue 82-003)* 2009; 20(2): 1-8.
53. Horsman J, Furlong W, Feeney D, et al. The Health Utilities Index (HUI): Concepts, measurement properties and applications. *Health and Quality of Life Outcomes* 2003; 1.
54. Kavirajan H, Hays CD, Vassar S, et al. Responsiveness and construct validity of the health utilities index in patients with dementia. *Medical Care* 2009; 47: 651-61.
55. Tuokko H, Woodward TS. Development and validation of a demographic correction system for neuropsychological measures used in the Canadian Study of Health and Aging. *Journal of Clinical and Experimental Neuropsychology* 1996; 18(4): 479-616.
56. Pierce JC, Cornell RG. Integrating stratum-specific likelihood ratios with the analysis of ROC curves. *Medical Decision Making* 1993; 13: 141-51.
57. Furukawa TA, Goldberg DP, Rabe-Hesketh S, et al. Stratum-specific likelihood ratios of two versions of the General Health Questionnaire. *Psychological Medicine* 2001; 31: 519-29.
58. Wada K, Tamaka K, Theriault G, et al. Application of the stratum-specific likelihood ratio (SSLR) analysis to results of a depressive symptoms screening survey among Japanese workers. *Society of Psychiatry and Psychiatric Epidemiology* 2007; 42: 410-3.
59. Schmitz N, Kruse J, Röss W. Application of stratum-specific likelihood ratios in mental health screening. *Social Psychiatry and Psychiatric Epidemiology* 2000; 35: 375-9.
60. Rust K, Rao JNK. Variance estimation for complex surveys using replication techniques. *Statistical Methods in Medical Research* 1996; 5: 281-310.
61. Backman L, Jones S, Small BJ, et al. Rate of cognitive decline in preclinical Alzheimer's disease: The role of comorbidity. *Journal of Gerontology: Psychological Sciences* 2003; 58B: P228-36.
62. Johnson KC, Margolis KL, Espeland MA, et al. A prospective study of the effect of hypertension and baseline blood pressure on cognitive decline and dementia in postmenopausal women: The women's health initiative memory study. *Journal of the American Geriatric Society* 2008; 56: 1449-58.
63. Peters R, Poulter R, Beckett N, et al. Cardiovascular and biochemical risk factors for incident dementia in the Hypertension in the Very Elderly Trial. *Journal of Hypertension* 2009; 27: 2055-62.
64. Muller M, Grobbee DE, Aleman A, et al. Cardiovascular disease and cognitive performance in middle-aged and elderly men. *Atherosclerosis* 2007; 190: 143-9.

## Appendix

**Table A**  
**Percentage distribution of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by delayed recall score and selected health characteristics, household population aged 45 or older, Canada excluding territories**

Health characteristic	Delayed recall t-score				
	Low 0 to 34	35 to 44	45 to 54	55 to 64	High 65 or more
	%				
<b>Total</b>	5.6	23.7	39.2	23.4	8.1
<b>Self-perceived health</b>					
Fair/Poor	5.8	29.1	40.6	19.2	5.2
Excellent/Very good/Good	5.6	22.8	39.0	24.1	8.5
<b>Self-perceived mental health</b>					
Fair/Poor	6.2 <sup>E</sup>	36.2	40.6	12.9	4.2 <sup>E</sup>
Excellent/Very good/Good	5.6	23.0	39.1	24.0	8.3
<b>Activities of daily living</b>					
No problems	6.6	31.4	38.1	18.0	5.9 <sup>F</sup>
Mild/Moderate/Severe/Total problems	5.5	23.0	39.3	23.9	8.3
<b>Life satisfaction</b>					
Low	8.1	27.8	40.3	18.4	5.4
Not low	5.2	22.9	39.0	24.4	8.6
<b>Depression</b>					
0.9 probability or higher	6.9 <sup>F</sup>	25.9	38.4	20.0	8.7 <sup>F</sup>
Less than 0.9 probability	5.5	23.5	39.3	23.7	8.1
<b>Loneliness</b>					
High	7.9	26.0	38.5	21.0	6.6
Not high	5.3	23.3	39.4	23.7	8.3
<b>Memory</b>					
Able to remember most things	5.4	22.9	39.3	23.8	8.6
Somewhat forgetful/Very forgetful/Unable to remember anything	6.3	26.3	38.8	22.3	6.4
<b>Ability to think clearly and solve problems</b>					
Able to think clearly/solve problems	5.3	23.4	39.2	23.8	8.3
Having a little/some/great deal of difficulty/unable	9.9	27.1	39.2	19.1	4.6 <sup>E</sup>
<b>Neurological disorder</b>					
Yes	9.8 <sup>F</sup>	32.1	39.0	16.6	F
No	5.5	23.5	39.2	23.6	8.2
<b>Vascular disorder</b>					
Yes	4.6	25.0	39.8	23.0	7.6
No	6.4	22.7	38.8	23.7	8.4
<b>Psychiatric disorder</b>					
Yes	7.3	26.9	39.4	20.1	6.3
No	5.4	23.3	39.2	23.8	8.3

<sup>E</sup> interpret with caution (coefficient of variation 16.6% to 33.3%)

<sup>F</sup> too unreliable to be reported (coefficient of variation greater than 33.3%)

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

**Table B**  
**Percentage distribution of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by animal-naming score and selected health characteristics, household population aged 45 or older, Canada excluding territories**

Health characteristic	Animal-naming t-score				
	Low 0 to 34	35 to 44	45 to 54	55 to 64	High 65 or more
	%				
<b>Total</b>	7.4	24.9	36.6	23.2	7.8
<b>Self-perceived health</b>					
Fair/Poor	9.0	30.0	37.3	18.4	5.3
Excellent/Very good/Good	7.1	24.0	36.5	24.0	8.3
<b>Self-perceived mental health</b>					
Fair/Poor	12.7	31.5	34.1	16.4	5.4 <sup>E</sup>
Excellent/Very good/Good	7.1	24.5	36.8	23.6	8.0
<b>Activities of daily living</b>					
No problems	7.4	24.3	36.5	23.8	8.0
Mild/Moderate/Severe/Total problems	7.6	31.2	37.5	18.0	5.7
<b>Life satisfaction</b>					
Low	12.4	30.3	32.8	18.1	6.3
Not low	6.6	23.9	37.3	24.2	8.1
<b>Depression</b>					
0.9 probability or higher	9.2 <sup>E</sup>	22.5	34.6	26.2	7.5
Less than 0.9 probability	7.3	25.0	36.7	23.1	7.8
<b>Loneliness</b>					
High	9.6	28.7	34.0	20.6	7.0
Not high	7.1	24.4	37.0	23.6	7.9
<b>Memory</b>					
Able to remember most things	6.7	24.4	37.1	23.5	8.3
Somewhat forgetful/Very forgetful/Unable to remember anything	9.7	26.6	35.2	22.4	6.2
<b>Ability to think clearly and solve problems</b>					
Able to think clearly/solve problems	6.9	24.5	36.7	23.8	8.1
Having a little/some/great deal of difficulty/unable	13.4	30.3	35.2	16.5	4.6 <sup>E</sup>
<b>Neurological disorder</b>					
Yes	12.2	26.9	39.2	16.7	5.1 <sup>E</sup>
No	7.3	24.9	36.5	23.4	7.9
<b>Vascular disorder</b>					
Yes	7.0	24.7	38.7	22.9	6.8
No	7.8	25.2	34.9	23.5	8.6
<b>Psychiatric disorder</b>					
Yes	8.3	27.3	37.3	21.9	5.2
No	7.3	24.7	36.5	23.4	8.1

<sup>E</sup> interpret with caution (coefficient of variation 16.6% to 33.3%)

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

*Validation of cognitive functioning categories in the Canadian Community Health Survey • Methodological insights*

**Table C**  
**Percentage distribution of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by Mental Alternation Test score and selected health characteristics, household population aged 45 or older, Canada excluding territories**

Health characteristic	Mental Alternation Test t-score				
	Low 0 to 34	35 to 44	45 to 54	55 to 64	High 65 or more
<b>Total</b>	6.5	23.2	36.4	26.0	7.9
<b>Self-perceived health</b>			%		
Fair/Poor	8.4	32.4	36.7	17.7	4.9
Excellent/Very good/Good	6.1	21.7	36.3	27.4	8.4
<b>Self-perceived mental health</b>					
Fair/Poor	10.0	32.8	32.1	20.3	4.8 <sup>E</sup>
Excellent/Very good/Good	6.2	22.7	36.6	26.4	8.1
<b>Activities of daily living</b>					
No problems	6.4	22.4	36.2	26.8	8.2
Mild/Moderate/Severe/Total problems	7.3	30.6	37.6	19.2	5.3
<b>Life satisfaction</b>					
Low	9.0	28.9	34.6	21.8	5.6
Not low	6.0	22.2	36.8	26.8	8.3
<b>Depression</b>					
0.9 probability or higher	6.9 <sup>E</sup>	25.1	34.7	28.1	5.2 <sup>E</sup>
Less than 0.9 probability	6.4	23.0	36.6	25.9	8.1
<b>Loneliness</b>					
High	8.6	26.6	36.1	22.7	6.0
Not high	6.2	22.7	36.5	26.4	8.2
<b>Memory</b>					
Able to remember most things	6.2	22.3	36.6	26.5	8.4
Somewhat forgetful/Very forgetful/Unable to remember anything	7.3	26.1	35.6	24.5	6.4
<b>Ability to think clearly and solve problems</b>					
Able to think clearly/solve problems	6.0	22.6	36.3	26.8	8.3
Having a little/some/great deal of difficulty/unable	11.4	30.8	36.8	17.5	3.6
<b>Neurological disorder</b>					
Yes	11.9 <sup>E</sup>	29.5	34.2	18.6	5.8 <sup>E</sup>
No	6.3	23.1	36.4	26.3	8.0
<b>Vascular disorder</b>					
Yes	6.4	24.7	37.2	23.6	8.2
No	6.5	22.0	35.8	28.0	7.7
<b>Psychiatric disorder</b>					
Yes	7.1	25.6	33.4	27.4	6.6 <sup>E</sup>
No	6.4	23.0	36.7	25.9	8.1

<sup>E</sup> interpret with caution (coefficient of variation 16.6% to 33.3%)

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

**Table D**  
**Stratum-specific likelihood ratios (SSLR) for selected health characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by delayed recall score category, household population aged 45 or older, Canada excluding territories**

Health characteristic	Delayed recall t-score														
	0 to 34			35 to 44			45 to 54			55 to 64			65 or more		
	SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval	
	from	to	from	to	from	to	from	to	from	to	from	to	from	to	
Low self-rated health	1.03	1.02	1.04	1.28	1.27	1.28	1.04	1.04	1.04	0.80	0.80	0.80	0.62	0.61	0.62
Low self-rated mental health	1.10	1.09	1.11	1.57	1.57	1.58	1.04	1.03	1.04	0.54	0.53	0.54	0.51	0.50	0.52
Difficulties with activities of daily living	1.19	1.18	1.20	1.37	1.36	1.37	0.97	0.97	0.97	0.75	0.75	0.75	0.71	0.71	0.72
Low life satisfaction	1.57	1.56	1.58	1.21	1.21	1.22	1.03	1.03	1.04	0.76	0.75	0.76	0.63	0.62	0.63
High probability of depression	1.09	1.08	1.10	1.11	1.10	1.11	0.96	0.96	0.97	0.87	0.86	0.87	1.20	1.19	1.21
High loneliness	1.49	1.48	1.50	1.12	1.11	1.12	0.98	0.98	0.98	0.88	0.88	0.89	0.79	0.79	0.80
Unable to remember things	1.16	1.15	1.16	1.15	1.14	1.15	0.99	0.98	0.99	0.94	0.94	0.94	0.74	0.74	0.75
Unable to think/solve problems	1.87	1.85	1.88	1.16	1.15	1.16	1.00	1.00	1.00	0.81	0.80	0.81	0.56	0.55	0.56
Neurological disorder	1.77	1.75	1.79	1.37	1.36	1.38	0.99	0.99	1.00	0.70	0.70	0.71	0.31	0.31	0.32
Vascular disorder	0.71	0.71	0.72	1.10	1.10	1.10	1.03	1.03	1.03	0.97	0.97	0.97	0.91	0.90	0.91
Psychiatric disorder	1.34	1.33	1.35	1.16	1.15	1.16	1.00	1.00	1.01	0.84	0.84	0.85	0.77	0.76	0.77

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

**Table E**  
**Stratum-specific likelihood ratios (SSLR) for selected health characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by animal-naming score category, household population aged 45 or older, Canada excluding territories**

Health characteristic	Animal-naming t-score														
	0 to 34			35 to 44			45 to 54			55 to 64			65 or more		
	SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval	
	from	to	from	to	from	to	from	to	from	to	from	to	from	to	
Low self-rated health	1.27	1.26	1.27	1.25	1.24	1.25	1.02	1.02	1.02	0.77	0.76	0.77	0.64	0.64	0.65
Low self-rated mental health	1.78	1.77	1.79	1.29	1.28	1.29	0.93	0.92	0.93	0.69	0.69	0.70	0.67	0.67	0.68
Difficulties with activities of daily living	1.02	1.01	1.03	1.28	1.28	1.29	1.03	1.03	1.03	0.76	0.75	0.76	0.71	0.71	0.72
Low life satisfaction	1.89	1.89	1.90	1.27	1.27	1.27	0.88	0.88	0.88	0.75	0.75	0.75	0.78	0.78	0.79
High probability of depression	1.20	1.19	1.20	0.87	0.87	0.87	0.98	0.98	0.98	1.16	1.16	1.17	0.86	0.85	0.86
High loneliness	1.36	1.35	1.37	1.18	1.17	1.18	0.92	0.92	0.92	0.87	0.87	0.88	0.89	0.88	0.89
Unable to remember things	1.43	1.43	1.44	1.09	1.09	1.09	0.95	0.95	0.95	0.95	0.95	0.96	0.75	0.74	0.75
Unable to think/solve problems	1.94	1.93	1.95	1.24	1.23	1.24	0.96	0.96	0.96	0.69	0.69	0.70	0.57	0.57	0.58
Neurological disorder	1.67	1.66	1.69	1.08	1.07	1.09	1.07	1.07	1.08	0.71	0.71	0.72	0.64	0.63	0.65
Vascular disorder	0.90	0.89	0.90	0.98	0.98	0.98	1.11	1.11	1.11	0.97	0.97	0.98	0.78	0.78	0.79
Psychiatric disorder	1.13	1.12	1.13	1.11	1.10	1.11	1.02	1.02	1.02	0.94	0.93	0.94	0.64	0.64	0.65

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

*Validation of cognitive functioning categories in the Canadian Community Health Survey • Methodological insights*

**Table F**  
**Stratum-specific likelihood ratios (SSLR) for selected health characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by Mental Alternation Test score category, household population aged 45 or older, Canada excluding territories**

Health characteristic	Mental Alternation Test t-score														
	0 to 34			35 to 44			45 to 54			55 to 64			65 or more		
	SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval		SSLR	95% confidence interval	
	from	to	from	to	from	to	from	to	from	to	from	to	from	to	
Low self-rated health	1.36	1.35	1.37	1.49	1.49	1.50	1.01	1.01	1.01	0.64	0.64	0.65	0.58	0.58	0.59
Low self-rated mental health	1.61	1.59	1.62	1.45	1.44	1.45	0.88	0.87	0.88	0.77	0.76	0.77	0.59	0.59	0.60
Difficulties with activities of daily living	1.15	1.15	1.16	1.36	1.36	1.37	1.04	1.03	1.04	0.72	0.71	0.72	0.65	0.64	0.65
Low life satisfaction	1.51	1.50	1.52	1.31	1.30	1.31	0.94	0.94	0.94	0.81	0.81	0.82	0.68	0.68	0.68
High probability of depression	1.11	1.10	1.12	1.04	1.04	1.05	0.97	0.97	0.98	1.08	1.08	1.09	0.65	0.64	0.66
High loneliness	1.40	1.39	1.41	1.17	1.17	1.17	0.99	0.99	0.99	0.86	0.86	0.86	0.73	0.72	0.73
Unable to remember things	1.18	1.17	1.18	1.17	1.17	1.17	0.97	0.97	0.98	0.93	0.92	0.93	0.77	0.76	0.77
Unable to think/solve problems	1.88	1.87	1.89	1.36	1.36	1.37	1.01	1.01	1.01	0.65	0.65	0.66	0.43	0.43	0.44
Neurological disorder	1.88	1.86	1.90	1.28	1.27	1.29	0.94	0.94	0.94	0.71	0.70	0.71	0.73	0.72	0.74
Vascular disorder	0.98	0.97	0.98	1.12	1.12	1.12	1.04	1.04	1.04	0.84	0.84	0.84	1.06	1.05	1.06
Psychiatric disorder	1.11	1.10	1.12	1.12	1.11	1.12	0.91	0.91	0.91	1.06	1.05	1.06	0.81	0.81	0.82

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

**Table G**  
**Odds ratios relating selected health status characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by delayed recall score, household population aged 45 or older, Canada excluding territories**

Health characteristic	Adjusted degrees of freedom	Adjusted chi-square	Delayed recall t-score											
			0 to 34			35 to 44			45 to 54			55 to 64		
			Odds ratio	95% confidence interval		Odds ratio	95% confidence interval		Odds ratio	95% confidence interval		Odds ratio	95% confidence interval	
	from	to	from	to	from	to	from	to	from	to	from	to		
Low self-rated health	3.89	28.86	1.67	1.10	2.53	2.07	1.44	2.98	1.69	1.21	2.35	1.30	0.91	1.83
Low self-rated mental health	3.43	38.94	2.16	1.24	3.74	3.08	1.91	4.98	2.03	1.29	3.21	1.05	0.65	1.69
Difficulties with activities of daily living	3.30	23.94	1.67	0.97	2.90	1.92	1.18	3.14	1.36	0.84	2.21	1.05	0.64	1.73
Low life satisfaction	3.71	29.47	2.51	1.64	3.82	1.93	1.40	2.67	1.65	1.24	2.20	1.21	0.89	1.65
High probability of depression	3.86	3.07	1.17	0.60	2.27	1.02	0.63	1.67	0.90	0.56	1.47	0.78	0.46	1.34
High loneliness	3.97	12.05	1.88	1.22	2.89	1.41	1.00	1.98	1.23	0.88	1.73	1.11	0.79	1.57
Unable to remember things	3.93	10.67	1.56	1.07	2.27	1.55	1.17	2.04	1.33	1.01	1.74	1.26	0.94	1.70
Unable to think/solve problems	3.77	18.96	3.36	1.72	6.58	2.09	1.15	3.80	1.80	0.98	3.30	1.45	0.76	2.75
Neurological disorder	3.77	20.40	5.65	2.34	13.66	4.38	1.97	9.72	3.17	1.43	7.06	2.25	0.98	5.14
Vascular disorder	3.93	12.07	0.79	0.58	1.07	1.21	0.96	1.53	1.13	0.92	1.40	1.07	0.86	1.32
Psychiatric disorder	3.88	9.67	1.75	1.07	2.84	1.51	1.04	2.19	1.31	0.91	1.88	1.10	0.75	1.61

Note: Comparison group is 65 or more t-score category.

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.



**Table H**  
**Odds ratios relating selected health status characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by animal-naming score, household population aged 45 or older, Canada excluding territories**

Health characteristic	Adjusted degrees of freedom	Adjusted chi-square	Animal-naming t-score											
			0 to 34			35 to 44			45 to 54			55 to 64		
			Odds ratio	95% confidence interval from to		Odds ratio	95% confidence interval from to		Odds ratio	95% confidence interval from to		Odds ratio	95% confidence interval from to	
Low self-rated health	3.82	39.18	1.97	1.42	2.75	1.95	1.50	2.53	1.59	1.24	2.04	1.19	0.91	1.57
Low self-rated mental health	3.62	27.76	2.64	1.59	4.38	1.91	1.19	3.06	1.38	0.88	2.14	1.03	0.69	1.55
Difficulties with activities of daily living	3.92	34.77	1.43	0.98	2.09	1.80	1.30	2.48	1.44	1.05	1.97	1.06	0.77	1.47
Low life satisfaction	3.70	57.94	2.42	1.70	3.43	1.62	1.23	2.14	1.12	0.85	1.48	0.96	0.71	1.28
High probability of depression	3.55	3.60	1.31	0.76	2.25	0.94	0.66	1.33	0.98	0.70	1.39	1.18	0.82	1.72
High loneliness	3.86	14.20	1.53	1.02	2.30	1.33	0.97	1.82	1.04	0.77	1.40	0.98	0.72	1.34
Unable to remember things	3.68	19.96	1.92	1.43	2.58	1.46	1.17	1.81	1.27	1.03	1.56	1.28	1.01	1.61
Unable to think/solve problems	3.67	36.37	3.38	2.09	5.47	2.16	1.44	3.24	1.67	1.12	2.50	1.21	0.77	1.90
Neurological disorder	3.78	13.17	2.61	1.20	5.66	1.68	0.84	3.37	1.67	0.85	3.30	1.11	0.51	2.41
Vascular disorder	3.90	13.50	1.14	0.87	1.50	1.25	1.03	1.52	1.41	1.17	1.71	1.24	1.02	1.52
Psychiatric disorder	3.68	7.87	1.75	1.16	2.65	1.72	1.25	2.36	1.59	1.18	2.13	1.46	1.06	2.00

Note: Comparison group is 65 or more t-score category.

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.

**Table I**  
**Odds ratios relating selected health status characteristics of respondents to 2009 Canadian Community Health Survey—Healthy Aging Cognition Module, by Mental Alternation Test score, household population aged 45 or older, Canada excluding territories**

Health characteristic	Adjusted degrees of freedom	Adjusted chi-square	Mental Alternation Test t-score											
			0 to 34			35 to 44			45 to 54			55 to 64		
			Odds ratio	95% confidence interval from to		Odds ratio	95% confidence interval from to		Odds ratio	95% confidence interval from to		Odds ratio	95% confidence interval from to	
Low self-rated health	3.83	99.35	2.34	1.67	3.28	2.56	1.95	3.37	1.74	1.31	2.30	1.11	0.84	1.46
Low self-rated mental health	3.78	26.20	2.71	1.55	4.74	2.44	1.38	4.32	1.48	0.86	2.55	1.30	0.74	2.27
Difficulties with activities of daily living	3.92	57.02	1.78	1.25	2.51	2.10	1.58	2.79	1.60	1.20	2.12	1.10	0.83	1.47
Low life satisfaction	3.89	42.25	2.23	1.54	3.23	1.92	1.44	2.57	1.39	1.06	1.81	1.20	0.90	1.59
High probability of depression	3.90	4.80	1.69	0.93	3.07	1.70	1.07	2.71	1.48	0.94	2.32	1.68	1.06	2.67
High loneliness	3.78	16.45	1.92	1.26	2.92	1.60	1.18	2.19	1.36	1.00	1.85	1.18	0.86	1.62
Unable to remember things	3.89	15.72	1.53	1.11	2.11	1.52	1.19	1.94	1.27	1.01	1.58	1.20	0.94	1.54
Unable to think/solve problems	3.59	47.03	4.35	2.67	7.09	3.15	2.15	4.60	2.34	1.61	3.39	1.51	1.00	2.29
Neurological disorder	3.87	21.92	2.58	1.28	5.22	1.76	0.98	3.16	1.29	0.70	2.36	0.97	0.51	1.83
Vascular disorder	3.95	17.98	0.92	0.71	1.20	1.06	0.87	1.30	0.98	0.81	1.18	0.80	0.65	0.97
Psychiatric disorder	3.93	4.96	1.37	0.83	2.25	1.37	0.90	2.09	1.12	0.75	1.68	1.30	0.86	1.96

Note: Comparison group is 65 or more t-score category.

Source: 2009 Canadian Community Health Survey—Healthy Aging Cognition Module.