

Article

Physical activity among First Nations people off reserve, Métis and Inuit

by Leanne C. Findlay

February 2011



Physical activity among First Nations people off reserve, Métis and Inuit

by *Leanne C. Findlay*

Abstract

Background

Research on physical activity among Aboriginal peoples has generally taken a pan-Aboriginal approach rather than considering First Nations people, Métis and Inuit separately. However, the groups differ geographically and culturally.

Data and methods

Data from the 2005 Canadian Community Health Survey were used to compare rates of active and moderately active leisure time (versus inactive) among First Nations people off reserve, Métis and Inuit with rates among non-Aboriginal people. Factors associated with active and moderately active leisure time were examined using logistic regression models. An active or moderately active lifestyle was studied in relation to self-perceived physical and mental health and the presence of chronic conditions.

Results

First Nations (people off reserve) and Métis people were significantly more likely than non-Aboriginal Canadians to have an active lifestyle. Being male, younger age and high educational attainment were associated with an increased likelihood of physically active leisure time. An active lifestyle was associated with an increased likelihood of excellent or very good self-perceived physical and mental health among Métis and among Aboriginal people overall. Level of leisure-time physical activity was not associated with chronic conditions for any Aboriginal group or for the non-Aboriginal population.

Interpretation

Aboriginal peoples generally have higher levels of leisure-time activity than do non-Aboriginal people.

Keywords

Aboriginal peoples, chronic disease, exercise, health status, leisure activities, mental health

Author

Leanne C. Findlay (613-951-4648; leanne.findlay@statcan.gc.ca) is with the Health Analysis Division at Statistics Canada in Ottawa, Ontario K1A 0T6.

Previous research has suggested that Aboriginal people off reserve may be more active than their non-Aboriginal counterparts.^{1,2} As well, according to results of a 2002/2003 survey, one-fifth of First Nations people living on reserves engaged in at least 30 minutes of moderate-to-vigorous activity four or more days per week. Whether they are Aboriginal^{3,4} or non-Aboriginal,² physically active people are more likely than those who are less active to report excellent or very good health.

Studies of physical activity among Aboriginal people have tended to consider First Nations people, Métis and Inuit collectively rather than as separate groups, or have focused exclusively on First Nations people. However, geographic and cultural differences between the groups may be related to leisure-time physical activity. Increasingly, the need for research that makes distinctions between First Nations people, Métis people and Inuit is being recognized.⁵

Relatively little information is available about factors that may influence Aboriginal peoples' participation in physical activity. A 2006 review⁶ found negative associations with age and body weight, and positive associations with education and perceived health. In addition, males were generally more

active than females, and people with a supportive social environment were more likely to be physically active. However, this review cited mostly American data; little is known about the correlates of physical activity for Aboriginal people in Canada, and specifically, separate Aboriginal groups.

The current study has three goals: 1) to examine leisure-time physical activity among First Nations people living off reserve, Métis people and Inuit; 2) to determine factors associated with active and moderately active (compared with inactive) leisure time for the three groups; and 3) to examine the relationship between physical activity and health among Aboriginal people.

Methods

Data source

Data from the 2005 Canadian Community Health Survey (CCHS) were used to examine the leisure-time physical activity of First Nations people off reserve, Métis and Inuit aged 12 or older. The target population of the CCHS is all Canadians aged 12 or older. Excluded from the sampling frame are individuals living on Indian Reserves and on Crown Lands, institutional residents, full-time members of the Canadian Forces, and residents of certain remote regions. Coverage is in the range of 98% in the provinces; in the Territories, it is about 90% in the Yukon, 97% in the Northwest Territories and 71% in Nunavut, primarily because some remote regions are excluded. In Nunavut, the CCHS collects information in the 10 largest communities: Iqaluit, Rankin Inlet, Cambridge Bay, Kugluktuk are always in sample, plus one community from Cape Dorset, Pangnirtung, Igloodik or Pond Inlet is selected; plus one community from Baker Lake or Arviat. Households were selected using a complex cluster design based on the Labour Force Survey.

CCHS respondents were asked, “Are you an Aboriginal person, that is, North American Indian, Métis or Inuit?” (Although respondents self-identified as “North American Indian,” the term “First Nations” is used throughout this study.) The 2005 sample included 3,414 respondents who self-identified as belonging to a single Aboriginal group (1,522 First Nations, 1,533 Métis, and 359 Inuit) and 129,494 respondents who were not Aboriginal. This analysis excludes 39 respondents who reported belonging to more than one Aboriginal group.

Measures

Respondents’ *leisure-time physical activity* was based on a list of common activities (Appendix Table A). Respondents reported the number of times they had engaged in each activity during the previous three months and the average duration. Average daily energy expenditure was calculated for each

activity by multiplying by an estimate of the energy cost of the activity (kilocalories per kilogram of body weight per hour, according to the Canadian Fitness and Lifestyle Research Institute guidelines). The energy expenditures were summed and used to classify respondents into one of three categories: active (3 or more kilocalories per kilogram of body weight per day); moderately active (1.5 to less than 3 kilocalories per kilogram per day); and inactive (less than 1.5 kilocalories per kilogram per day). Active leisure time is the equivalent of walking at least 1 hour every day; moderately active leisure time, half an hour per day.

Three health indicators were considered in the current study: *self-perceived health* (general); *self-perceived mental health*; and the presence of *chronic conditions* (for example, asthma, high blood pressure, arthritis) (Appendix Table B). Respondents were asked, “In general, would you say your [mental] health is...” The response options were: excellent, very good, good, fair, and poor. The number of chronic conditions a respondent reported was dichotomized to reflect the presence of one or more.

Several socio-demographic characteristics were examined as predictors of leisure-time physical activity: gender, age, number of dependants aged 0 to 17 in the household (asked of respondents aged 18 or older), marital status (married/common-law versus single/widowed/divorced/separated), and employment (yes or no). Age was classified into five groups: 12 to 17, 18 to 34, 35 to 49, 50 to 64, and 65 or older. Total annual household income was classified into three categories: less than \$20,000; \$20,000 to \$39,999; and \$40,000 or more. Education was classified as: less than secondary graduation, secondary graduation, some postsecondary, and postsecondary graduation.

Statistical analyses

Descriptive statistics on the socio-demographic characteristics, self-reported health indicators, and leisure-time physical activity of the study sample were calculated. To account for

the younger age profile of the Aboriginal population, percentages for all health outcomes and leisure-time physical activities were age-standardized to the Aboriginal population. Chi-square comparisons were used to identify significant differences between each Aboriginal group and the non-Aboriginal population (but not between Aboriginal groups). Logistic regression was used to determine whether the socio-demographic factors were associated with active and/or moderately active leisure time, and to examine associations between the level of leisure-time activity and the measures of self-reported health. Separate models were fitted for each Aboriginal group and for the non-Aboriginal comparison group. Because of the relatively small sample for each Aboriginal group, models combining the three were also fitted. Sampling weights were used in all analyses. To account for the complex survey design, a bootstrapping technique was applied for variance estimation.⁷

Results

More active/Less healthy

First Nations people who lived off-reserve and Métis people were more likely than the non-Aboriginal population to be physically active in their leisure time: 37% and 39% versus 30% (Table 1). However, the percentage of Inuit who were physically active (31%) was not significantly different from the percentage for non-Aboriginal people.

Each Aboriginal group was more likely than the non-Aboriginal population to report their general and mental health as good/fair/poor rather than excellent/very good. As well, the prevalence of chronic conditions was higher among First Nations people off reserve (71%) and Métis (74%) people than among non-Aboriginal Canadians (64%). The comparatively low rate (65%) among Inuit may be associated with the CCHS question, which specified chronic conditions “diagnosed by a health professional.” Inuit communities may have relatively few such people to make the diagnoses.

Table 1
Physical activity, health and socio-demographic characteristics, by Aboriginal identity, household population aged 12 or older, Canada, 2005

| Characteristics | First Nations off reserve (n=1,522) | | | Métis (n=1,533) | | | Inuit (n=359) | | | Non-Aboriginal (n=129,494) | | |
|---|-------------------------------------|-------|-----|-----------------|-------|-----|-------------------|-------|-----|----------------------------|------|-----|
| | % | Mean | se | % | Mean | se | % | Mean | se | % | Mean | se |
| Physical activity | | | | | | | | | | | | |
| Active | 37.0* | ... | ... | 38.5* | ... | ... | 30.7 | ... | ... | 29.9 | ... | ... |
| Moderately active | 22.1 | ... | ... | 21.8 | ... | ... | 24.5 | ... | ... | 25.0 | ... | ... |
| Inactive | 40.9 | ... | ... | 39.7 | ... | ... | 44.9 | ... | ... | 45.2 | ... | ... |
| Health | | | | | | | | | | | | |
| Self-rated health | | | | | | | | | | | | |
| Good/Fair/Poor | 48.3* | ... | ... | 45.8* | ... | ... | 51.3* | ... | ... | 36.2 | ... | ... |
| Excellent/Very good | 51.7 | ... | ... | 54.2 | ... | ... | 48.7 | ... | ... | 63.8 | ... | ... |
| Self-rated mental health | | | | | | | | | | | | |
| Good/Fair/Poor | 32.7* | ... | ... | 30.0* | ... | ... | 38.2* | ... | ... | 24.5 | ... | ... |
| Excellent/Very good | 67.3 | ... | ... | 70.0 | ... | ... | 61.8 | ... | ... | 75.5 | ... | ... |
| Chronic condition | | | | | | | | | | | | |
| No | 28.8* | ... | ... | 26.1* | ... | ... | 34.9 | ... | ... | 35.9 | ... | ... |
| Yes | 71.2 | ... | ... | 73.9 | ... | ... | 65.1 | ... | ... | 64.1 | ... | ... |
| Socio-demographic | | | | | | | | | | | | |
| Sex | | | | | | | | | | | | |
| Male | 51.9 | ... | ... | 50.6 | ... | ... | 56.4 | ... | ... | 49.3 | ... | ... |
| Female | 48.1 | ... | ... | 49.4 | ... | ... | 43.6 | ... | ... | 50.8 | ... | ... |
| Age | | | | | | | | | | | | |
| | ... | 35.2* | 0.6 | ... | 36.6* | 0.7 | ... | 33.3* | 1.5 | ... | 43.0 | 0.0 |
| Marital status | | | | | | | | | | | | |
| Married/Common-law | 47.0* | ... | ... | 52.8* | ... | ... | 44.3* | ... | ... | 59.0 | ... | ... |
| Single/Widowed/Divorced/Separated | 53.1 | ... | ... | 47.2 | ... | ... | 55.7 | ... | ... | 41.0 | ... | ... |
| Number of children in household† | | | | | | | | | | | | |
| | ... | 0.9* | 0.1 | ... | 0.8* | 0.1 | ... | 1.3* | 0.2 | ... | 0.6 | 0.0 |
| Education | | | | | | | | | | | | |
| Less than secondary graduation | 36.2* | ... | ... | 36.0* | ... | ... | 54.1* | ... | ... | 23.7 | ... | ... |
| Secondary graduation | 15.5 | ... | ... | 14.8 | ... | ... | 7.1 ^E | ... | ... | 15.2 | ... | ... |
| Some postsecondary | 12.5 | ... | ... | 10.0 | ... | ... | 7.1 ^E | ... | ... | 8.8 | ... | ... |
| Postsecondary graduation | 35.9 | ... | ... | 39.3 | ... | ... | 31.8 ^E | ... | ... | 52.4 | ... | ... |
| Employment | | | | | | | | | | | | |
| Currently working | 65.2* | ... | ... | 68.4 | ... | ... | 60.1 | ... | ... | 69.2 | ... | ... |
| Not working | 34.8 | ... | ... | 31.6 | ... | ... | 39.9 | ... | ... | 30.9 | ... | ... |
| Household income | | | | | | | | | | | | |
| Less than \$20,000 | 18.5* | ... | ... | 14.5* | ... | ... | 18.0 ^E | ... | ... | 9.4 | ... | ... |
| \$20,000 to \$39,999 | 25.8 | ... | ... | 21.2 | ... | ... | 23.6 ^E | ... | ... | 18.9 | ... | ... |
| More than \$40,000 | 55.8 | ... | ... | 64.3 | ... | ... | 58.4 | ... | ... | 71.7 | ... | ... |

* significantly different distribution from non-Aboriginal population (p<0.05)

† population aged 18 or older

se = standard error

... not applicable

Note: Chi-square comparisons have been age-standardized to Aboriginal population.

Source: 2005 Canadian Community Health Survey.

To some extent, these differences in physical activity and health reflect socio-demographic characteristics. Each Aboriginal group tended to be younger and to have more dependent children in their households than did non-Aboriginal

Canadians. They were also more likely to live in rural areas, to be single, to live in households with low annual income, and to have less than secondary graduation. First Nations people off reserve were also significantly less likely to be employed.

Active leisure time

The first set of models examined socio-demographic characteristics that might be related to active (versus inactive) leisure time (Table 2). For First Nations people off reserve and Métis people, gender, education and age were significantly associated with active leisure time. The odds of being active were higher for males than for females. Individuals with higher levels of education were more likely to be active than were those who had not graduated from secondary school. And 12- to 17-year-olds had significantly higher odds of active leisure time than did people aged 18 to 34. The odds of active leisure time were significantly lower among First Nations people off reserve aged 35 to 64. For Inuit, no socio-demographic factors were significantly related to active leisure time, although the small sample size may have precluded the detection of significant associations.

To put the findings for each Aboriginal group in context, a model was fitted for non-Aboriginal Canadians. Most of the socio-demographic characteristics included in the model were significantly related to leisure-time physical activity. The odds of being active (versus inactive) were significantly higher for non-Aboriginal people who were male, whose education had advanced beyond secondary graduation, and who were aged 12 to 17. The odds of being active were significantly lower for those who lived in lower-income households (less than \$40,000), who were married or in a common-law relationship, who had a relatively large number of young dependants in their household, who were employed, and who were aged 35 or older.

Moderately active leisure time

Fewer characteristics were associated with moderately active (versus inactive) leisure time (Table 3). For First Nations people off reserve, household income below \$40,000 (rather than \$40,000 or more) was associated with lower odds of moderately active leisure time. For Métis people, urban residence and

Table 2
Odds ratios relating active (versus inactive) leisure time to selected characteristics, by Aboriginal identity, household population aged 12 or older, Canada, 2005

| | First Nations off reserve (n=1,522) | | | Métis (n=1,533) | | | Inuit (n=359) | | | Non-Aboriginal (n=129,494) | | |
|--|---|-------------------------------|-------|--------------------|-------------------------------|-------|------------------|-------------------------------|--------|-------------------------------|-------------------------------|------|
| | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | |
| | | from | to | | from | to | | from | to | | from | to |
| Sex | | | | | | | | | | | | |
| Male | 1.66* | 1.07 | 2.59 | 1.72* | 1.08 | 2.75 | 3.54 | 0.85 | 14.76 | 1.35* | 1.28 | 1.42 |
| Female† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Age group | | | | | | | | | | | | |
| 12 to 17 | 3.97* | 1.24 | 12.68 | 3.88* | 1.09 | 13.74 | 6.42 | 0.61 | 67.95 | 2.86* | 2.50 | 3.27 |
| 18 to 34† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| 35 to 49 | 0.53* | 0.30 | 0.92 | 1.01 | 0.55 | 1.85 | 0.23 | 0.03 | 1.73 | 0.72* | 0.68 | 0.76 |
| 50 to 64 | 0.38* | 0.18 | 0.81 | 0.88 | 0.40 | 1.90 | 0.16 | 0.00 | 14.21 | 0.67* | 0.63 | 0.72 |
| 65 or older | 0.40 | 0.15 | 1.09 | 0.71 | 0.21 | 2.41 | 0.26 | 0.00 | 609.41 | 0.72* | 0.65 | 0.79 |
| Marital status | | | | | | | | | | | | |
| Married/Common-law | 0.89 | 0.54 | 1.47 | 0.58 | 0.31 | 1.12 | 1.06 | 0.26 | 4.25 | 0.73* | 0.69 | 0.77 |
| Single/Widowed/Divorced/Separated† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Number of children in household | | | | | | | | | | | | |
| | 0.96 | 0.78 | 1.18 | 1.20 | 0.92 | 1.57 | 1.17 | 0.68 | 2.01 | 0.96* | 0.93 | 0.99 |
| Education | | | | | | | | | | | | |
| Less than secondary graduation† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Secondary graduation | 0.82 | 0.40 | 1.70 | 2.00 | 0.81 | 4.94 | 0.62 | 0.02 | 19.24 | 1.41* | 1.29 | 1.55 |
| Some postsecondary | 2.80* | 1.24 | 6.31 | 2.27 | 0.78 | 6.55 | 2.72 | 0.15 | 48.60 | 1.57* | 1.40 | 1.76 |
| Postsecondary graduation | 2.01* | 1.13 | 3.59 | 3.82* | 1.81 | 8.10 | 3.78 | 0.52 | 27.41 | 1.75* | 1.62 | 1.89 |
| Employment | | | | | | | | | | | | |
| Working | 0.60 | 0.34 | 1.05 | 0.69 | 0.36 | 1.33 | 0.63 | 0.12 | 3.26 | 0.74* | 0.69 | 0.79 |
| Not working† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Household income | | | | | | | | | | | | |
| Less than \$20,000 | 0.69 | 0.34 | 1.37 | 0.61 | 0.28 | 1.36 | 1.01 | 0.13 | 7.74 | 0.61* | 0.57 | 0.67 |
| \$20,000 to \$39,999 | 0.72 | 0.42 | 1.25 | 0.69 | 0.35 | 1.34 | 1.50 | 0.18 | 12.47 | 0.65* | 0.61 | 0.69 |
| More than \$40,000† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |

† reference category

* significantly different from estimate for reference category ($p < 0.05$)

... not applicable

Note: All models control for population density.

Source: 2005 Canadian Community Health Survey.

postsecondary graduation were associated with increased odds of moderate activity. None of the socio-demographic factors was associated with moderate activity among Inuit. However, among the non-Aboriginal population, being older, lower education and household income, being employed and being married were negatively associated with moderately active leisure time.

Leisure-time activity and self-perceived health

To determine if active or moderately active leisure time was associated with health status over and above socio-demographic characteristics, separate

models were fitted for each Aboriginal group, for the three groups combined, and for the non-Aboriginal population.

Active leisure time was associated with self-perceived general and mental health for Métis people (Table 4). The estimates were not significant for First Nations people off reserve or Inuit, although the odds ratios were in the same direction. To overcome the small sample sizes, the three groups were considered together. In this case, active leisure time was associated with increased self-perceived general and mental health, even when controlling for socio-demographic characteristics. For non-Aboriginal Canadians both active and moderately

active leisure time were associated with increased odds of reporting excellent or very good self-perceived health.

Neither active nor moderately active leisure time was associated with the presence of one or more chronic conditions for any Aboriginal group (alone or combined) or for the non-Aboriginal population.

Discussion

According to results of the 2005 Canadian Community Health Survey, First Nations people living off-reserve and Métis people were significantly more physically active in their leisure time

Table 3
Odds ratio relating moderately active leisure time (versus inactive) to selected characteristics, by Aboriginal identity, household population aged 12 or older, Canada, 2005

| | First Nations off reserve (n=1,522) | | | Métis (n=1,533) | | | Inuit (n=359) | | | Non-Aboriginal (n=129,494) | | |
|--|-------------------------------------|------------------------------------|------|-----------------|------------------------------------|------|---------------|------------------------------------|---------|----------------------------|------------------------------------|------|
| | Odds ratio | 95% confidence interval from to | | Odds ratio | 95% confidence interval from to | | Odds ratio | 95% confidence interval from to | | Odds ratio | 95% confidence interval from to | |
| Sex | | | | | | | | | | | | |
| Male | 1.00 | 0.59 | 1.68 | 1.03 | 0.64 | 1.65 | 1.01 | 0.26 | 3.90 | 1.03 | 0.98 | 1.08 |
| Female† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Age group | | | | | | | | | | | | |
| 12 to 17 | 1.86 | 0.43 | 7.96 | 1.26 | 0.35 | 4.57 | 0.61 | 0.00 | 1008.97 | 1.73* | 1.49 | 2.01 |
| 18 to 34† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| 35 to 49 | 0.63 | 0.34 | 1.20 | 0.78 | 0.44 | 1.39 | 0.47 | 0.10 | 2.28 | 0.90* | 0.84 | 0.96 |
| 50 to 64 | 0.81 | 0.39 | 1.67 | 0.76 | 0.38 | 1.53 | 1.45 | 0.27 | 7.85 | 0.91* | 0.85 | 0.98 |
| 65 or older | 1.76 | 0.63 | 4.93 | 1.38 | 0.44 | 4.34 | 0.35 | 0.00 | 2466.37 | 1.08 | 0.98 | 1.18 |
| Marital status | | | | | | | | | | | | |
| Married/Common-law | 1.11 | 0.63 | 1.96 | 0.82 | 0.45 | 1.49 | 0.83 | 0.24 | 2.86 | 0.92* | 0.87 | 0.97 |
| Single/Widowed/Divorced/Separated† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Number of children in household | 1.03 | 0.82 | 1.31 | 0.97 | 0.76 | 1.24 | 1.19 | 0.68 | 2.06 | 0.96* | 0.93 | 0.99 |
| Education | | | | | | | | | | | | |
| Less than secondary graduation† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Secondary graduation | 1.68 | 0.77 | 3.63 | 1.24 | 0.55 | 2.79 | 0.90 | 0.04 | 22.43 | 1.33* | 1.22 | 1.46 |
| Some postsecondary | 1.29 | 0.52 | 3.15 | 0.89 | 0.36 | 2.23 | 1.10 | 0.19 | 6.31 | 1.48* | 1.33 | 1.64 |
| Postsecondary graduation | 1.16 | 0.58 | 2.32 | 2.12* | 1.15 | 3.89 | 1.46 | 0.36 | 5.86 | 1.63* | 1.51 | 1.75 |
| Employment | | | | | | | | | | | | |
| Working | 0.96 | 0.51 | 1.83 | 1.07 | 0.57 | 1.99 | 0.83 | 0.25 | 2.78 | 0.83* | 0.77 | 0.88 |
| Not working† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Household income | | | | | | | | | | | | |
| Less than \$20,000 | 0.48* | 0.23 | 0.97 | 0.67 | 0.30 | 1.54 | 1.24 | 0.26 | 6.06 | 0.64* | 0.59 | 0.69 |
| \$20,000 to \$39,999 | 0.33* | 0.18 | 0.62 | 0.90 | 0.49 | 1.65 | 1.03 | 0.21 | 5.01 | 0.72* | 0.68 | 0.77 |
| More than \$40,000† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |

† reference category

* significantly different from estimate for reference category (p<0.05)

... not applicable

Note: All models control for population density.

Source: 2005 Canadian Community Health Survey.

than was the non-Aboriginal population. These results reinforce earlier findings,^{1,2} although in this analysis, significant differences were not apparent for Inuit.

Many of the factors that influence leisure-time activity in the general population were significant for Aboriginal peoples.^{2,6,8} Being male, younger age and higher educational attainment were associated with *active* leisure among First Nations people off reserve and Métis people. Fewer factors were associated with *moderately active* leisure among Aboriginal peoples (possibly because of small sample sizes), despite significant relationships for the non-Aboriginal population.

As reported in earlier studies,²⁻⁴ whether they were Aboriginal or non-Aboriginal, physically active people were more likely than those who were less active to report excellent or very good general and mental health. However, moderately active leisure time was not positively associated with self-perceived general and mental health for Aboriginal people, which indicates a need for further research to determine the amount of activity required for associations to emerge.

Despite higher levels of physical activity, Aboriginal people tend to report poorer health. Earlier studies have shown a relatively high prevalence of obesity^{8,9,10}

and the associated complications of diabetes¹¹⁻¹³ and other chronic conditions³ among the Aboriginal population.

It is often suggested that health disparities are driven by social rather than biological determinants.¹⁴ For instance, smoking, poor housing conditions and lower income can negatively affect health.¹⁵ When the influence of several social determinants was taken into account, active leisure time was associated with health only for Métis people.

A more complex relationship between physical activity and health may exist for Aboriginal people. It is possible that factors not considered in this analysis

Table 4

Adjusted odds ratios relating level of leisure time activity to self-rated general and mental health and presence of chronic condition(s), by Aboriginal identity, household population aged 12 or older, Canada, 2005

| | First Nations off reserve (n=1,522) | | | Métis (n=1,533) | | | Inuit (n=359) | | | Total Aboriginal (n=3,414) | | | Non-Aboriginal (n=129,494) | | |
|---------------------------------------|-------------------------------------|-------------------------|------|-----------------|-------------------------|------|---------------|-------------------------|-------|----------------------------|-------------------------|------|----------------------------|-------------------------|------|
| | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | | Odds ratio | 95% confidence interval | |
| | | from | to | | from | to | | from | to | | from | to | | from | to |
| Self-rated health | | | | | | | | | | | | | | | |
| Active leisure time | 1.50 | 0.93 | 2.41 | 3.48* | 2.04 | 5.93 | 2.13 | 0.66 | 6.92 | 2.24* | 1.59 | 3.17 | 2.16* | 2.04 | 2.29 |
| Moderately active leisure time | 0.93 | 0.56 | 1.52 | 1.55 | 0.93 | 2.56 | 1.34 | 0.36 | 4.97 | 1.22 | 0.88 | 1.70 | 1.52* | 1.44 | 1.61 |
| Inactive leisure time† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| Self-rated mental health | | | | | | | | | | | | | | | |
| Active leisure time | 1.55 | 0.98 | 2.46 | 1.74* | 1.00 | 3.03 | 0.62 | 0.18 | 2.18 | 1.58* | 1.12 | 2.22 | 1.52* | 1.43 | 1.62 |
| Moderately active leisure time | 1.47 | 0.90 | 2.40 | 1.12 | 0.67 | 1.85 | 1.69 | 0.61 | 4.71 | 1.31 | 0.94 | 1.81 | 1.29* | 1.21 | 1.36 |
| Inactive leisure time† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |
| One or more chronic conditions | | | | | | | | | | | | | | | |
| Active leisure time | 1.62 | 0.93 | 2.81 | 0.82 | 0.47 | 1.44 | 3.39 | 0.92 | 12.44 | 1.24 | 0.84 | 1.82 | 0.93 | 0.87 | 0.98 |
| Moderately active leisure time | 1.70 | 0.95 | 3.05 | 0.69 | 0.39 | 1.20 | 1.38 | 0.46 | 4.15 | 1.12 | 0.76 | 1.64 | 1.04 | 0.98 | 1.10 |
| Inactive leisure time† | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... | 1.00 | ... | ... |

† reference category

* significantly different from estimate for reference category ($p < 0.05$)

... not applicable

Note: All models control for age, sex, marital status, number of dependant children in household, education, employment status, household income, and population density.

Source: 2005 Canadian Community Health Survey.

What is already known on this subject?

- Aboriginal peoples in Canada have relatively high rates of obesity and diabetes.
- Active leisure time is associated with better health.
- Little information is available for the three Aboriginal groups—First Nations people off reserve, Métis, Inuit—separately.

What does this study add?

- First Nations people off reserve and Métis are more active than non-Aboriginal Canadians.
- Socio-demographic characteristics are associated with physically active leisure time.
- Active leisure time is associated with better self-perceived general and mental health for Aboriginal people.

may be influential. The well-being of Aboriginal people may have not only physical and mental components, but also emotional and spiritual aspects based on a holistic approach.^{16,17} Wilson et al.¹⁸ suggested that in addition to well-established social determinants (such as income and education), traditional activities may be significantly associated with the health of Aboriginal people.

In the current study, active and moderately active leisure time were not significantly related to the presence of chronic conditions. However, the data are cross-sectional, and it is not possible to determine temporal associations. In addition, the total number of conditions and the the severity of impairment were not considered. By contrast, previous research has demonstrated links between physical activity and specific chronic disease risk factors. For instance, Katzmarzyk⁹ found an association between Aboriginal identity and obesity, and between physical activity and obesity. Physical activity has also been shown to be inversely associated with metabolic syndrome among men

(which, in turn, is associated with an elevated risk for cardiovascular disease and diabetes).¹⁹ Links between physical activity and specific chronic conditions among Aboriginal people warrant further investigation.

Strengths, limitations and future directions

The current study provides insight into the levels of leisure-time activity among a population-based sample of First Nations people off reserve, Métis people and the Inuit. Previous work in this area has not focused on factors associated with physical activity, or on associations between physical activity and the health of specific Aboriginal groups.

Some limitations should be acknowledged. Because the data are cross-sectional, it is not possible to determine the direction of relationships between physical activity and health in this study. Healthier people may be capable of active leisure time, rather than active leisure time leading to better health.

The specific activities in which Aboriginal people participated could not be identified. Moreover, the CCHS measure of physical activity may not be suited to Aboriginal people. It is based on a predetermined list of “common” activities that may not be prevalent in Aboriginal communities, while activities relevant to Aboriginal lifestyles are not included. For example, results from the First Nations Regional Longitudinal Health Survey revealed that hunting, trapping, berry-picking and food gathering (none of which were listed in the CCHS) were frequently cited as physical activities by First Nations people.⁴ As well, self-rated health may not be construed in the same way by Aboriginal people because of a more holistic perspective¹⁶ or because the CCHS categories are understood differently.

Similarly, the thresholds for *active*, *moderately active* and *inactive* leisure time may not be appropriate for Aboriginal people. They may not consider some of the activities listed by the CCHS as leisure, and therefore, do not include them among their leisure-time pursuits.²⁰ Kriska et al.²¹ found that occupational activity was a greater contributor to total physical activity than was leisure time among Aboriginal people. If this is, indeed, the case, levels of physical activity may be underestimated in the current study. Additional research is necessary to determine if the measure of physical activity in the CCHS is appropriate for Aboriginal people.

Although First Nations people off reserve, Métis people and Inuit were examined separately, the leisure-time activities of even smaller groups with diverse traditions, history, and culture¹⁰ might warrant attention. However,

given that the current study was already limited by sample size, such research is not feasible using the CCHS. In fact, the failure to detect significant differences for factors potentially associated with leisure-time activity or between an active or moderately active lifestyle and health may be the result of small sample sizes.

Conclusion

The current study demonstrates that First Nations people off reserve and Métis people were more likely than non-Aboriginal Canadians to have an active lifestyle. The analysis highlights the importance of examining the three Aboriginal groups separately and the value of studying the relationship between physical activity and health. With an estimate of close to 1.2 million Aboriginal people in Canada in 2006,²² further research is needed to identify factors associated with their physical and mental well-being. ■

References

- Bryan SN, Tremblay MS, Perez CE, et al. Physical activity and ethnicity: Evidence from the Canadian Community Health Survey. *Canadian Journal of Public Health* 2006; 97(4): 271-6.
- Gilmour H. Physically active Canadians. *Health Reports* (Statistics Canada, Catalogue 82-003) 2007; 18(3): 45-65.
- Janz T, Seto J, Turner A. *Aboriginal Peoples Survey, 2006: An Overview of the Health of the Métis Population* (Statistics Canada, Catalogue 89-637-C, No. 004) Ottawa: Minister of Industry, 2009.
- First Nations Information Governance Committee. *First Nations Regional Longitudinal Health Survey 2002/03*. Ottawa: Assembly of First Nations/First Nations Information Governance Committee, 2007.
- Task Force on Aboriginal Languages and Cultures. *Towards a New Beginning: A Foundational Report for a Strategy to Revitalize First Nations, Metis and Inuit languages and Cultures*. Ottawa: Ministry of Canadian Heritage, 2005.
- Coble JD, Rhodes RE. Physical activity and Native Americans. *American Journal of Preventative Medicine* 2006; 31(1): 36-46.
- Rust K, Rao JNK. Variance estimation for complex surveys using replication techniques. *Statistical Methods in Medical Research* 1996; 5: 281-310.
- Bryan S, Walsh P. Physical activity and obesity in Canadian women. *BMC Women's Health* 2004; 4(suppl 1): S6.
- Katzmarzyk PT. Obesity and physical activity among Aboriginal Canadians. *Obesity* 2007; 16(1): 184-90.
- Tremblay MS, Perez CE, Ardern CI et al. Obesity, overweight and ethnicity. *Health Reports* (Statistics Canada, Catalogue 82-003) 2005; 16(4): 23-34.
- Adelson N. The embodiment of inequity: Health disparities in Aboriginal Canada. *Canadian Journal of Public Health* 2005; 96: S45-62.
- Bruce S. The impact of diabetes mellitus among the Métis of western Canada. *Ethnicity and Health* 2000; 5(1): 47-57.
- Young TK, Reading J, Elias B, O'Neil JD. Type 2 diabetes mellitus in Canada's First Nations: Status of an epidemic in progress. *Canadian Medical Association Journal* 2000; 163(5): 561-6.
- Smylie J. *Indigenous Children's Health Report: Health Assessment in Action*. Toronto: Keenan Research Centre, 2009.
- Tjepkema, M. The health of the off-reserve Aboriginal population. *Health Reports* (Statistics Canada, Catalogue 82-003) 2002; 13: 1-17.
- King M, Smith A, Gray-Donald K. Indigenous health part 2: The underlying causes of the health gap. *Lancet* 2009; 374: 76-85.
- Anderson M, Smylie J, Anderson I, et al. *First Nations, Métis, and Inuit Health Indicators in Canada: A Background Paper for the Project "Action-oriented indicators for health and healthy systems development for indigenous peoples in Australia, Canada, and New Zealand."* Discussion Paper No. 18. Melbourne, Australia: School of Population Health, University of Melbourne, 2006.
- Wilson K, Rosenberg MW. Exploring the determinants of health for First Nations peoples in Canada: can existing frameworks accommodate traditional activities? *Social Science and Medicine* 2002; 55: 2017-31.
- Liu J, Young TK, Zinman B, et al. Lifestyle variables, non-traditional cardiovascular risk factors, and the metabolic syndrome in an Aboriginal Canadian population. *Obesity* 2006; 14(3): 500-98.
- Thompson SJ, Gifford SM, Thorpe L. The social and cultural context of risk and prevention: Food and physical activity in an urban Aboriginal community. *Health Education and Behavior* 2000; 27(6): 725-43.
- Kriska AM, Hanley AJG, Harris SB, Zinman B. Physical activity, physical fitness, and insulin and glucose concentrations in an isolated native Canadian population experiencing rapid lifestyle change. *Diabetes Care* 2001; 24(10): 1787-92.
- Statistics Canada. *Aboriginal Peoples in Canada in 2006: Inuit, Métis and First Nations, 2006 Census* (Catalogue 97-558-XIE) Ottawa: Minister of Industry, 2008.

Appendix

Table A
Selected activities and metabolic equivalent of task (MET) value, 2005 Canadian Community Health Survey

| Activity | MET value (kilocalories per kilogram per hour) |
|-------------------------------|---|
| Jogging/Running | 9.5 |
| Basketball | 6.0 |
| Ice hockey | 6.0 |
| In-line skating/Rollerblading | 5.0 |
| Soccer | 5.0 |
| Volleyball | 5.0 |
| Bicycling | 4.0 |
| Downhill skiing/Snowboarding | 4.0 |
| Exercise class/Aerobics | 4.0 |
| Golfing | 4.0 |
| Ice skating | 4.0 |
| Tennis | 4.0 |
| Baseball/Softball | 3.0 |
| Fishing | 3.0 |
| Gardening/Yard work | 3.0 |
| Home exercises | 3.0 |
| Popular or social dance | 3.0 |
| Swimming | 3.0 |
| Walking for exercise | 3.0 |
| Weight training | 3.0 |
| Bowling | 2.0 |
| Other* | 4.0 |

* mean MET value applied for all "other" activities

Table B
Chronic conditions listed in 2005 Canadian Community Health Survey

| Condition |
|--|
| Food allergies |
| Other allergies |
| Asthma |
| Fibromyalgia |
| Arthritis/Rheumatism |
| Back problems |
| High blood pressure |
| Migraine headaches |
| Chronic bronchitis |
| Emphysema |
| Chronic obstructive pulmonary disease |
| Diabetes |
| Epilepsy |
| Heart disease |
| Cancer |
| Intestinal/Stomach ulcers |
| Effects of stroke |
| Urinary incontinence |
| Bowel disorder |
| Alzheimer's disease/Other dementia |
| Cataracts |
| Glaucoma |
| Thyroid condition |
| Chronic fatigue syndrome |
| Multiple chemical sensitivities |
| Schizophrenia |
| Mood disorder such as depression, bipolar disorder, mania, dysthymia |
| Anxiety disorder such as a phobia, obsessive compulsive disorder, panic disorder |
| Autism/Other developmental disorder |
| Learning disability |
| Eating disorder (anorexia, bulimia) |
| Other long-term physical or mental health condition diagnosed by health professional |