

Table 4-2

Major salivary glands — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Major salivary glands (ICD-O-3: C07.9-C08.9) ³									
1992/1994									
Canada	67	62	71	62	56	68	73	67	78
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	67	61	72	62	55	69	74	65	81
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	62	46	76	F	F	F	F	F	F
British Columbia	72	60	82	67	49	81	77	62	88
1993/1995									
Canada	68	64	71	62	56	68	75	69	80
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	70	65	75	66	59	73	75	66	82
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	63	52	73	F	F	F	F	F	F
British Columbia	71	60	80	66	51	79	79	65	89
1994/1996									
Canada	66	62	69	59	53	64	75	69	80
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	65	60	70	59	52	66	74	65	81
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	71	62	78	60	47	71	81	69	90
1995/1997									
Canada	65	61	68	59	53	64	73	67	78
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	65	60	70	59	52	66	74	65	82
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	66	57	74	F	F	F	74	63	84
1996/1998									
Canada	66	62	70	59	54	64	75	70	81
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	66	60	71	58	51	65	78	69	86
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	73	61	83	F	F	F	F	F	F
British Columbia	69	60	77	F	F	F	78	67	86

See footnotes at the end of the table.

Table 4-2 – continued

Major salivary glands — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
1997/1999									
Canada	69	65	73	64	58	69	78	73	83
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	72	66	76	65	58	72	83	74	90
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	69	56	80	F	F	F	F	F	F
British Columbia	68	58	76	60	44	73	79	67	88
1998/2000									
Canada	68	65	72	61	56	66	79	73	84
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	70	65	74	65	58	71	80	71	87
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	66	53	78	F	F	F	F	F	F
British Columbia	69	60	77	F	F	F	86	74	94
1999/2001									
Canada	69	65	73	64	58	69	77	71	82
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	69	64	73	66	60	72	72	64	79
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	69	56	80	F	F	F	F	F	F
British Columbia	71	62	79	61	47	73	F	F	F
2000/2002									
Canada	69	66	73	65	60	70	75	69	80
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	70	65	74	68	62	74	72	64	78
Manitoba	81	61	96	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	68	54	79	F	F	F	F	F	F
British Columbia	66	56	75	56	46	66	F	F	F
2001/2003									
Canada	73	70	77	70	65	75	77	71	82
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	74	69	78	72	65	77	76	69	83
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	79	65	89	F	F	F	F	F	F
British Columbia	73	62	82	64	51	74	F	F	F

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1572).