

Table 4-3

Major salivary glands — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Major salivary glands (ICD-O-3: C07.9-C08.9) ³									
Newfoundland and Labrador									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	69 ^E	34 ^E	98 ^E	58 ^E	14 ^E	112 ^E	F	F	F
1999/2001	82 ^E	47 ^E	108 ^E	80 ^E	33 ^E	120 ^E	F	F	F
2000/2002	85 ^E	54 ^E	108 ^E	83 ^E	41 ^E	116 ^E	F	F	F
2001/2003	84 ^E	55 ^E	101 ^E	84 ^E	48 ^E	104 ^E	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Nova Scotia									
1992/1994	67	44	85	59	30	85	79	40	97
1993/1995	58	38	77	43	19	69	83	48	100
1994/1996	59	39	76	42	20	65	90	52	103
1995/1997	54	33	73	47	23	70	68	29	94
1996/1998	55	30	76	54	25	78	F	F	F
1997/1999	65	37	86	60	28	84	F	F	F
1998/2000	59	32	82	49	18	78	F	F	F
1999/2001	70	43	91	63	29	91	78	37	104
2000/2002	65	39	86	66	30	93	64	27	91
2001/2003	65	38	87	61	26	90	70	29	96
New Brunswick									
1992/1994	57	28	82	F	F	F	F	F	F
1993/1995	38	18	60	14	1	51	63	29	87
1994/1996	58	35	78	32	8	64	82	47	98
1995/1997	63	39	81	43	16	71	87	47	101
1996/1998	77	53	93	69	36	91	88	48	103
1997/1999	64	40	81	55	27	79	76	36	97
1998/2000	57	35	75	44	21	67	F	F	F
1999/2001	65	42	83	46	21	71	97	51	107
2000/2002	62	38	80	44	18	69	F	F	F
2001/2003	77	53	92	74	43	94	F	F	F
Ontario									
1992/1994	67	60	73	63	54	70	73	63	82
1993/1995	69	63	75	65	56	73	76	66	84
1994/1996	65	59	71	59	51	67	74	65	82
1995/1997	65	59	71	58	50	66	75	66	83
1996/1998	65	59	71	57	49	65	78	69	86
1997/1999	71	65	76	63	55	70	83	74	90
1998/2000	69	63	74	61	53	68	81	72	88
1999/2001	68	62	73	63	56	71	75	66	82
2000/2002	69	64	74	66	58	73	73	66	80
2001/2003	74	68	78	69	62	76	78	71	84

See footnotes at the end of the table.

Table 4-3 – continued

Major salivary glands — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	53	34	71	31	13	54	91	55	103
1993/1995	63	41	81	51	26	75	86	44	102
1994/1996	70	46	87	58	26	85	82	47	98
1995/1997	71	46	88	F	F	F	62	31	84
1996/1998	61	37	82	60	18	100	62	33	84
1997/1999	54	29	78	55	17	95	54	22	79
1998/2000	63	40	84	58	26	91	72	37	93
1999/2001	66	42	86	70	34	97	63	29	86
2000/2002	76	54	93	82	54	102	63	29	89
2001/2003	65	43	83	76	44	99	51	23	77
Saskatchewan									
1992/1994	66	46	84	68	31	98	65	40	84
1993/1995	75	49	94	75	31	110	73	42	93
1994/1996	78	49	99	80	37	113	75	38	100
1995/1997	64	38	87	55	23	86	75	35	103
1996/1998	56	37	75	40	18	64	76	44	99
1997/1999	59	38	78	53	27	77	69	34	94
1998/2000	61	42	78	47	22	70	78	48	96
1999/2001	64	44	81	56	30	80	77	44	95
2000/2002	63	44	80	48	25	72	82	52	99
2001/2003	69	49	86	55	28	80	86	55	102
Alberta									
1992/1994	67	51	80	68	44	86	66	44	82
1993/1995	66	50	78	63	42	79	70	46	87
1994/1996	64	48	77	62	40	80	66	42	84
1995/1997	69	54	81	61	41	78	79	55	94
1996/1998	73	58	85	65	40	86	79	59	92
1997/1999	74	59	86	62	38	81	85	65	96
1998/2000	70	55	83	51	30	71	90	70	99
1999/2001	71	56	82	50	30	68	92	73	101
2000/2002	68	53	81	48	27	68	86	65	98
2001/2003	76	60	89	71	47	89	81	58	96
British Columbia									
1992/1994	70	58	80	65	48	80	75	58	88
1993/1995	68	56	78	62	47	76	76	58	89
1994/1996	69	59	78	59	44	72	80	66	90
1995/1997	66	56	75	56	41	69	75	61	86
1996/1998	70	59	78	58	42	72	79	65	89
1997/1999	68	57	77	57	41	71	79	63	90
1998/2000	69	58	78	53	39	67	86	70	96
1999/2001	72	61	81	57	42	70	90	74	99
2000/2002	69	58	79	52	37	66	90	74	99
2001/2003	75	64	83	62	48	75	92	77	100

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).