

Table 7-3

Nasopharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Nasopharynx (ICD-O-3: C11.0-C11.9) ³									
Newfoundland and Labrador									
1992/1994	28 ^E	7 ^E	56 ^E	33 ^E	8 ^E	63 ^E	F	F	F
1993/1995	46 ^E	17 ^E	74 ^E	41 ^E	13 ^E	71 ^E	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	49 ^E	21 ^E	75 ^E	40 ^E	12 ^E	68 ^E	F	F	F
1999/2001	43 ^E	13 ^E	72 ^E	43 ^E	13 ^E	72 ^E	F	F	F
2000/2002	43 ^E	13 ^E	72 ^E	43 ^E	13 ^E	72 ^E	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Nova Scotia									
1992/1994	45	22	66	49	20	73	F	F	F
1993/1995	46	21	69	53	22	78	F	F	F
1994/1996	49	22	72	67	31	89	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	44	16	70	F	F	F	F	F	F
1997/1999	47	17	74	F	F	F	F	F	F
1998/2000	43	20	66	48	20	73	F	F	F
1999/2001	41	15	66	42	13	70	F	F	F
2000/2002	32	8	61	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
New Brunswick									
1992/1994	67	35	91	55	23	84	F	F	F
1993/1995	68	36	93	60	27	87	F	F	F
1994/1996	56	27	80	59	27	86	F	F	F
1995/1997	41	15	66	48	18	75	F	F	F
1996/1998	42	13	70	F	F	F	F	F	F
1997/1999	63	27	87	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	62	26	86	F	F	F	F	F	F
Ontario									
1992/1994	61	54	67	56	48	64	72	59	82
1993/1995	63	56	69	60	51	67	71	58	81
1994/1996	63	56	69	61	52	68	67	55	77
1995/1997	62	56	69	61	53	69	64	52	74
1996/1998	64	57	70	64	55	71	64	51	75
1997/1999	65	58	71	65	57	72	65	53	75
1998/2000	64	58	70	64	56	71	65	53	75
1999/2001	65	59	70	64	56	71	67	55	76
2000/2002	62	56	68	59	51	66	70	58	79
2001/2003	64	57	70	61	53	68	71	58	81

See footnotes at the end of the table.

Table 7-3 – continued

Nasopharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
	percent								
Manitoba									
1992/1994	54	29	75	58	30	79	F	F	F
1993/1995	53	28	75	57	28	80	F	F	F
1994/1996	51	29	71	57	30	79	F	F	F
1995/1997	34	16	54	38	17	60	F	F	F
1996/1998	31	13	52	27	8	52	F	F	F
1997/1999	49	25	70	51	25	74	F	F	F
1998/2000	69	43	87	83	47	99	F	F	F
1999/2001	70	45	87	80	48	97	F	F	F
2000/2002	53	31	72	46	17	73	60	28	82
2001/2003	53	26	76	F	F	F	F	F	F
Saskatchewan									
1992/1994	47	17	73	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	29	7	57	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	49	20	74	58	24	83	F	F	F
2000/2002	27	8	51	36	11	63	F	F	F
2001/2003	34	14	56	45	16	71	F	F	F
Alberta									
1992/1994	44	30	58	45	30	60	39	9	75
1993/1995	43	28	57	46	30	61	25	4	60
1994/1996	54	39	67	59	42	73	32	8	62
1995/1997	60	45	74	67	47	81	45	19	70
1996/1998	69	52	81	67	46	82	74	42	91
1997/1999	65	48	78	60	39	76	74	44	90
1998/2000	69	51	81	60	38	77	85	52	97
1999/2001	64	47	77	56	37	72	85	49	98
2000/2002	61	46	74	48	32	63	113	..	..
2001/2003	56	42	68	48	33	61	94	51	109
British Columbia									
1992/1994	61	51	70	59	45	70	65	48	78
1993/1995	60	50	69	58	45	69	63	46	76
1994/1996	64	54	74	62	48	74	69	52	81
1995/1997	70	60	78	69	57	79	71	53	83
1996/1998	72	63	80	70	58	79	78	61	89
1997/1999	73	64	81	70	59	79	81	63	91
1998/2000	74	65	81	71	60	80	79	64	89
1999/2001	71	62	78	68	57	77	77	61	87
2000/2002	69	60	77	66	54	75	76	61	87
2001/2003	69	60	77	63	52	73	82	65	92

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).