

Table 10-2

Other buccal cavity and pharynx — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Other buccal cavity and pharynx (ICD-O-3: C09.0-C09.9, C14.0, C14.2-C14.8) ³									
1992/1994									
Canada	46	43	50	43	38	47	56	49	63
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	49	44	54	48	42	54	54	44	63
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	49	41	58	44	33	54	F	F	F
1993/1995									
Canada	47	43	50	43	38	47	58	51	64
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	52	47	57	50	44	56	58	48	66
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	35	22	48	F	F	F	F	F	F
British Columbia	48	39	56	42	31	52	F	F	F
1994/1996									
Canada	46	42	49	43	39	48	51	44	58
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	48	43	53	46	40	52	55	45	64
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	33	22	44	F	F	F	F	F	F
British Columbia	48	39	56	47	37	56	F	F	F
1995/1997									
Canada	45	41	48	44	39	48	48	41	55
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	46	41	51	44	38	50	55	45	63
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	41	29	52	F	F	F	F	F	F
British Columbia	46	37	54	49	39	58	F	F	F
1996/1998									
Canada	46	43	50	46	42	50	48	42	55
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	46	41	51	46	40	52	47	38	56
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	47	36	58	F	F	F	F	F	F
British Columbia	45	37	54	48	39	58	F	F	F

See footnotes at the end of the table.

Table 10-2 – continued

Other buccal cavity and pharynx — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
1997/1999									
Canada	47	44	51	47	43	51	49	42	55
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	49	44	53	51	45	57	45	36	54
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	50	39	60	50	37	61	F	F	F
British Columbia	46	37	55	49	38	58	F	F	F
1998/2000									
Canada	50	46	53	50	46	53	50	43	56
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	42	30	54	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	53	49	58	56	51	62	47	38	56
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	42	31	52	41	30	52	F	F	F
British Columbia	45	37	53	43	34	53	F	F	F
1999/2001									
Canada	53	49	56	52	48	56	53	46	59
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	40	27	53	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	57	52	61	58	53	63	53	44	62
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	44	35	54	43	32	53	F	F	F
British Columbia	50	42	57	48	39	57	F	F	F
2000/2002									
Canada	54	50	57	53	49	57	55	49	61
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	59	54	63	59	54	64	58	49	67
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	48	38	57	46	34	57	F	F	F
British Columbia	47	40	55	45	37	54	F	F	F
2001/2003									
Canada	57	54	60	57	53	60	59	53	65
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	..	..	..
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	62	57	66	61	56	66	63	54	71
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	53	44	61	50	39	60	F	F	F
British Columbia	52	45	59	51	43	59	F	F	F

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1572).