

Table 10-3

Other buccal cavity and pharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Other buccal cavity and pharynx (ICD-O-3: C09.0-C09.9, C14.0, C14.2-C14.8) ³									
Newfoundland and Labrador									
1992/1994	32 ^E	12 ^E	57 ^E	33 ^E	10 ^E	62 ^E	F	F	F
1993/1995	32 ^E	12 ^E	57 ^E	33 ^E	10 ^E	61 ^E	F	F	F
1994/1996	40 ^E	15 ^E	67 ^E	F	F	F	F	F	F
1995/1997	27 ^E	6 ^E	54 ^E	F	F	F	F	F	F
1996/1998	45 ^E	20 ^E	69 ^E	60 ^E	25 ^E	86 ^E	F	F	F
1997/1999	40 ^E	18 ^E	63 ^E	39 ^E	14 ^E	66 ^E	F	F	F
1998/2000	52 ^E	29 ^E	72 ^E	55 ^E	28 ^E	77 ^E	F	F	F
1999/2001	57 ^E	33 ^E	76 ^E	53 ^E	24 ^E	76 ^E	F	F	F
2000/2002	63 ^E	38 ^E	81 ^E	61 ^E	32 ^E	83 ^E	F	F	F
2001/2003	63 ^E	40 ^E	80 ^E	54 ^E	28 ^E	75 ^E	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	..	..	..
Nova Scotia									
1992/1994	40	22	58	31	13	53	57	21	86
1993/1995	38	20	57	27	10	48	F	F	F
1994/1996	57	36	75	57	31	77	F	F	F
1995/1997	39	23	56	40	21	59	F	F	F
1996/1998	40	25	56	42	24	60	37	11	65
1997/1999	35	22	49	32	18	48	45	17	72
1998/2000	40	27	53	37	23	52	50	21	76
1999/2001	39	26	52	39	25	53	42	13	71
2000/2002	47	33	60	46	31	61	49	20	73
2001/2003	57	42	70	57	40	72	55	26	78
New Brunswick									
1992/1994	49	28	69	46	22	68	F	F	F
1993/1995	48	27	67	42	20	63	F	F	F
1994/1996	49	29	67	39	20	59	F	F	F
1995/1997	40	22	58	38	18	58	F	F	F
1996/1998	48	31	65	36	17	56	72	37	93
1997/1999	45	28	62	40	20	61	54	23	80
1998/2000	57	40	72	54	34	72	64	29	87
1999/2001	60	43	74	60	42	76	56	21	84
2000/2002	59	42	73	58	40	73	F	F	F
2001/2003	56	39	70	55	37	70	F	F	F
Ontario									
1992/1994	47	42	52	46	40	52	50	40	59
1993/1995	50	44	55	48	42	54	54	44	63
1994/1996	47	41	52	45	39	51	50	40	59
1995/1997	45	40	50	43	37	49	48	39	57
1996/1998	45	40	50	46	40	52	41	32	50
1997/1999	47	42	52	51	44	57	40	32	49
1998/2000	52	47	57	57	51	62	43	34	52
1999/2001	57	52	61	60	54	65	49	40	58
2000/2002	59	54	63	59	54	65	57	47	65
2001/2003	62	57	66	62	56	67	62	53	71

See footnotes at the end of the table.

Table 10-3 – continued

Other buccal cavity and pharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	40	24	57	33	16	53	56	23	82
1993/1995	34	18	51	22	7	43	54	23	80
1994/1996	35	20	52	30	13	50	46	17	73
1995/1997	44	28	59	41	22	60	49	22	72
1996/1998	51	35	66	50	30	68	53	24	76
1997/1999	54	38	68	50	30	67	63	33	84
1998/2000	53	37	68	44	26	62	76	41	96
1999/2001	57	40	71	48	30	65	88	45	104
2000/2002	54	38	68	58	39	74	46	21	69
2001/2003	58	41	72	65	45	81	40	15	66
Saskatchewan									
1992/1994	26	11	44	11	2	31	F	F	F
1993/1995	19	8	34	9	2	26	42	13	71
1994/1996	23	10	40	20	6	40	32	8	61
1995/1997	27	13	43	20	6	39	40	12	69
1996/1998	37	20	55	25	9	46	60	25	85
1997/1999	36	20	53	22	7	43	58	27	82
1998/2000	34	18	51	25	9	45	53	19	80
1999/2001	29	14	46	30	12	51	27	6	54
2000/2002	34	18	51	42	21	62	F	F	F
2001/2003	46	28	63	57	35	76	F	F	F
Alberta									
1992/1994	34	23	47	33	20	47	37	15	62
1993/1995	31	20	44	30	17	44	36	13	62
1994/1996	33	21	44	33	21	47	29	9	56
1995/1997	39	27	51	41	27	54	33	10	59
1996/1998	49	36	60	48	34	61	50	24	72
1997/1999	51	39	62	51	37	63	49	24	71
1998/2000	42	31	53	43	31	55	39	17	61
1999/2001	46	36	56	46	34	57	48	27	66
2000/2002	51	40	60	50	38	61	52	32	70
2001/2003	55	46	64	54	43	65	58	39	74
British Columbia									
1992/1994	47	37	56	41	30	52	60	42	75
1993/1995	47	38	56	40	29	50	65	47	78
1994/1996	46	37	54	45	34	55	49	33	63
1995/1997	43	35	51	46	36	56	36	23	50
1996/1998	43	34	51	47	37	57	31	18	45
1997/1999	45	36	53	48	38	58	35	20	52
1998/2000	44	36	53	43	33	53	47	30	64
1999/2001	49	41	57	49	40	58	50	34	66
2000/2002	48	40	56	46	37	55	54	38	68
2001/2003	52	44	59	53	44	61	49	34	64

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).