

Table 12-1

Stomach — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	1 year of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Stomach (ICD-O-3: C16.0-C16.9) ²									
1992									
Total, 15 to 99 years	20	18	22	18	16	21	22	18	25
15 to 44 years	26	18	35	18	9	30	34	22	47
45 to 54 years	24	17	30	24	16	32	23	13	35
55 to 64 years	21	16	25	21	16	26	20	13	29
65 to 74 years	20	17	24	18	14	23	24	18	31
75 to 99 years	17	13	20	16	11	21	18	13	24
1993									
Total, 15 to 99 years	19	17	21	19	16	21	18	15	22
15 to 44 years	31	22	40	32	20	45	29	17	42
45 to 54 years	19	13	25	15	10	22	27	16	41
55 to 64 years	22	17	27	23	18	28	19	11	28
65 to 74 years	18	15	22	19	15	23	18	13	24
75 to 99 years	16	12	19	16	11	21	15	11	21
1994									
Total, 15 to 99 years	21	19	23	20	17	22	22	19	26
15 to 44 years	34	24	44	38	24	51	29	15	44
45 to 54 years	29	23	36	26	19	34	36	24	48
55 to 64 years	20	16	24	19	14	24	23	16	31
65 to 74 years	22	19	26	19	15	24	28	22	34
75 to 99 years	15	12	19	16	12	22	14	10	20
1995									
Total, 15 to 99 years	20	18	22	17	15	19	25	22	29
15 to 44 years	22	15	31	24	14	36	20	9	33
45 to 54 years	27	21	33	25	18	32	31	20	41
55 to 64 years	19	15	23	15	11	20	28	20	37
65 to 74 years	20	17	24	17	13	21	26	20	32
75 to 99 years	18	15	22	15	11	20	23	17	30
1996									
Total, 15 to 99 years	22	20	24	20	18	23	24	21	28
15 to 44 years	23	15	31	23	14	33	22	12	35
45 to 54 years	24	18	30	23	17	31	26	15	39
55 to 64 years	25	20	30	24	19	30	27	19	35
65 to 74 years	24	21	28	21	17	26	30	23	37
75 to 99 years	18	15	21	16	12	21	20	15	25
1997									
Total, 15 to 99 years	21	19	23	18	16	20	27	23	30
15 to 44 years	28	20	37	21	12	32	39	24	54
45 to 54 years	24	18	31	23	16	31	26	16	38
55 to 64 years	19	15	24	17	13	23	24	16	34
65 to 74 years	22	19	26	19	15	23	29	23	36
75 to 99 years	19	16	23	16	12	21	24	18	30
1998									
Total, 15 to 99 years	22	20	24	21	18	23	24	21	28
15 to 44 years	35	26	44	31	20	43	39	25	52
45 to 54 years	26	20	32	24	17	31	29	19	41
55 to 64 years	27	22	32	23	18	29	36	27	46
65 to 74 years	21	17	24	20	16	25	22	16	28
75 to 99 years	18	15	22	18	13	23	19	14	25
1999									
Total, 15 to 99 years	21	19	23	20	17	22	23	20	26
15 to 44 years	30	21	39	31	18	45	29	18	41
45 to 54 years	31	24	38	31	23	39	32	20	44
55 to 64 years	20	16	24	18	14	23	24	16	34
65 to 74 years	22	19	26	20	17	25	27	20	34
75 to 99 years	16	13	19	15	11	20	17	12	22

See footnotes at the end of the table.

Table 12-1 – continued

Stomach — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	1 year of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
	percent								
2000									
Total, 15 to 99 years	21	19	23	21	19	24	22	19	25
15 to 44 years	18	11	26	13	6	24	24	13	37
45 to 54 years	25	20	31	21	15	28	33	23	44
55 to 64 years	26	21	31	26	20	31	25	17	35
65 to 74 years	22	19	26	22	18	27	22	16	28
75 to 99 years	18	15	22	18	14	23	18	14	24
2001									
Total, 15 to 99 years	22	20	24	21	19	24	23	20	26
15 to 44 years	27	18	36	28	16	40	25	14	39
45 to 54 years	29	23	35	29	21	36	30	20	41
55 to 64 years	26	22	31	24	19	30	32	23	42
65 to 74 years	22	19	26	19	15	24	29	23	37
75 to 99 years	16	13	20	18	14	23	14	10	19
2002									
Total, 15 to 99 years	25	22	27	23	20	26	27	24	31
15 to 44 years	37	27	46	43	29	57	30	18	43
45 to 54 years	23	17	29	21	14	28	28	18	40
55 to 64 years	28	24	33	24	19	30	36	27	45
65 to 74 years	28	24	33	28	23	33	30	23	37
75 to 99 years	19	15	22	17	13	22	22	16	27
2003									
Total, 15 to 99 years	24	22	26	24	22	27	25	21	28
15 to 44 years	38	29	47	37	26	48	40	26	53
45 to 54 years	30	24	36	34	27	42	23	15	33
55 to 64 years	25	21	30	25	20	30	26	18	35
65 to 74 years	27	23	31	26	21	31	30	23	39
75 to 99 years	18	15	22	17	13	22	20	15	25

See footnotes at the end of the table.

Table 12-1 – continued

Stomach — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Stomach (ICD-O-3: C16.0-C16.9) ²									
1992/1994									
Total, 15 to 99 years	20	18	21	19	18	20	21	19	23
15 to 44 years	30	25	35	29	22	36	31	23	39
45 to 54 years	24	21	28	22	18	26	29	22	36
55 to 64 years	21	18	23	21	18	24	21	17	26
65 to 74 years	20	18	22	19	16	21	23	20	27
75 to 99 years	16	14	18	16	13	19	16	13	19
1993/1995									
Total, 15 to 99 years	20	19	21	18	17	20	22	20	24
15 to 44 years	29	24	34	31	24	39	26	19	34
45 to 54 years	25	22	29	22	18	27	31	25	38
55 to 64 years	20	18	23	19	16	22	24	19	29
65 to 74 years	20	18	22	18	16	21	24	20	28
75 to 99 years	16	14	19	16	13	18	17	14	21
1994/1996									
Total, 15 to 99 years	21	20	22	19	18	20	24	22	26
15 to 44 years	26	21	31	27	21	34	23	16	31
45 to 54 years	27	23	30	25	21	29	31	24	38
55 to 64 years	21	19	24	19	16	22	26	21	31
65 to 74 years	22	20	24	19	17	22	28	24	32
75 to 99 years	17	15	19	16	13	19	19	16	22
1995/1997									
Total, 15 to 99 years	21	20	22	19	17	20	25	23	27
15 to 44 years	24	20	29	23	17	29	27	20	35
45 to 54 years	25	22	29	24	20	28	28	22	35
55 to 64 years	21	19	24	19	16	22	26	22	32
65 to 74 years	22	20	24	19	17	22	28	25	32
75 to 99 years	19	17	21	16	13	18	22	19	26
1996/1998									
Total, 15 to 99 years	22	20	23	20	18	21	25	23	27
15 to 44 years	28	23	33	25	19	31	34	26	42
45 to 54 years	25	21	28	23	19	28	27	21	34
55 to 64 years	24	21	27	22	19	25	29	24	34
65 to 74 years	22	20	24	20	18	23	27	23	31
75 to 99 years	18	16	21	16	14	19	21	18	24
1997/1999									
Total, 15 to 99 years	21	20	22	19	18	21	25	23	27
15 to 44 years	31	26	36	27	20	34	35	28	43
45 to 54 years	27	23	31	26	22	30	29	23	36
55 to 64 years	22	20	25	20	17	23	29	23	34
65 to 74 years	22	20	24	20	18	22	26	22	30
75 to 99 years	18	16	20	16	14	19	20	17	23
1998/2000									
Total, 15 to 99 years	21	20	23	20	19	22	23	21	25
15 to 44 years	28	23	33	25	18	32	31	24	38
45 to 54 years	27	24	31	25	21	29	31	25	38
55 to 64 years	24	22	27	22	19	26	29	24	34
65 to 74 years	22	20	24	21	19	24	23	20	27
75 to 99 years	17	15	19	17	14	20	18	15	21
1999/2001									
Total, 15 to 99 years	21	20	22	21	19	22	23	21	24
15 to 44 years	25	20	30	23	17	30	26	20	34
45 to 54 years	28	25	32	27	22	31	32	25	38
55 to 64 years	24	21	27	23	20	26	27	22	33
65 to 74 years	22	20	24	21	18	23	26	22	30
75 to 99 years	17	15	19	17	15	20	16	14	19

See footnotes at the end of the table.

Table 12-1 – continued

Stomach — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
2000/2002									
Total, 15 to 99 years	22	21	24	22	20	23	24	22	26
15 to 44 years	27	22	32	28	21	35	27	20	34
45 to 54 years	26	23	29	24	20	28	31	25	37
55 to 64 years	27	24	30	25	22	28	32	26	37
65 to 74 years	24	22	26	23	20	26	27	23	31
75 to 99 years	18	16	20	18	15	20	18	15	21
2001/2003									
Total, 15 to 99 years	24	22	25	23	21	24	25	23	27
15 to 44 years	34	29	39	36	29	43	32	24	40
45 to 54 years	28	24	31	28	24	33	27	21	33
55 to 64 years	27	24	29	24	21	28	32	27	37
65 to 74 years	26	24	28	24	22	27	30	26	34
75 to 99 years	18	16	20	17	15	20	18	15	21

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM tables 103-1569 and 103-1570).