

Table 13-3

Small intestine — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Small intestine (ICD-O-3: C17.0-C17.9) ³									
Newfoundland and Labrador									
1992/1994	42 ^E	15 ^E	68 ^E	F	F	F	F	F	F
1993/1995	41 ^E	19 ^E	64 ^E	36 ^E	11 ^E	64 ^E	F	F	F
1994/1996	59 ^E	34 ^E	80 ^E	54 ^E	22 ^E	81 ^E	F	F	F
1995/1997	57 ^E	31 ^E	79 ^E	64 ^E	27 ^E	91 ^E	F	F	F
1996/1998	54 ^E	29 ^E	76 ^E	63 ^E	27 ^E	91 ^E	44 ^E	14 ^E	74 ^E
1997/1999	43 ^E	23 ^E	62 ^E	53 ^E	19 ^E	80 ^E	36 ^E	13 ^E	61 ^E
1998/2000	41 ^E	22 ^E	60 ^E	F	F	F	45 ^E	20 ^E	68 ^E
1999/2001	44 ^E	22 ^E	65 ^E	46 ^E	14 ^E	76 ^E	43 ^E	16 ^E	69 ^E
2000/2002	37 ^E	15 ^E	60 ^E	48 ^E	15 ^E	80 ^E	F	F	F
2001/2003	25 ^E	9 ^E	47 ^E	32 ^E	10 ^E	59 ^E	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	58	21	87	F	F	F	F	F	F
1997/1999	51	21	77	F	F	F	F	F	F
1998/2000	47	21	72	45	16	74	F	F	F
1999/2001	34	10	62	39	12	68	F	F	F
2000/2002	28	7	57	25	4	59	F	F	F
2001/2003	28	7	56	F	F	F	F	F	F
Nova Scotia									
1992/1994	39	21	58	41	17	69	36	13	63
1993/1995	36	18	56	27	7	57	44	18	71
1994/1996	46	27	65	44	18	70	49	22	73
1995/1997	58	37	76	45	19	71	73	39	92
1996/1998	57	39	72	52	30	72	66	35	87
1997/1999	44	29	59	40	21	60	50	26	72
1998/2000	34	22	48	35	19	53	33	15	55
1999/2001	32	20	46	29	14	47	36	17	57
2000/2002	33	20	47	28	13	45	41	19	62
2001/2003	32	18	46	23	9	40	48	22	71
New Brunswick									
1992/1994	38	20	58	36	13	65	40	15	68
1993/1995	42	22	64	40	16	68	47	15	82
1994/1996	39	21	59	40	18	64	F	F	F
1995/1997	42	22	63	35	13	61	54	20	84
1996/1998	29	13	48	21	5	45	40	12	70
1997/1999	37	19	55	20	5	43	57	26	82
1998/2000	44	24	63	48	20	75	39	14	65
1999/2001	53	32	72	75	40	100	34	12	58
2000/2002	47	27	66	95	52	115	21	6	41
2001/2003	41	23	60	64	29	93	26	10	48
Ontario									
1992/1994	34	29	40	34	27	41	35	27	43
1993/1995	38	33	44	37	29	44	41	32	49
1994/1996	38	33	43	32	26	40	45	37	53
1995/1997	40	34	45	33	26	40	48	40	56
1996/1998	39	34	44	34	28	41	44	36	51
1997/1999	40	35	45	37	30	44	43	36	50
1998/2000	46	41	51	46	39	53	46	39	53
1999/2001	48	43	53	48	41	55	48	40	55
2000/2002	47	42	52	46	39	53	48	40	55
2001/2003	48	43	53	47	40	54	49	42	55

See footnotes at the end of the table.

Table 13-3 – continued

Small intestine — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	28	15	42	34	16	54	20	6	41
1993/1995	29	16	43	34	17	52	22	7	44
1994/1996	36	22	52	44	25	64	24	7	47
1995/1997	38	23	53	41	21	61	33	14	56
1996/1998	38	24	54	36	18	56	42	19	65
1997/1999	38	24	52	35	17	55	41	22	60
1998/2000	39	26	53	45	24	65	35	18	52
1999/2001	41	28	55	46	23	69	38	22	55
2000/2002	46	31	60	56	33	76	37	19	56
2001/2003	53	37	67	60	37	79	46	25	65
Saskatchewan									
1992/1994	51	34	67	58	33	80	42	20	64
1993/1995	49	31	66	45	24	66	57	26	84
1994/1996	48	31	65	51	29	71	43	18	70
1995/1997	47	29	65	46	23	69	49	22	74
1996/1998	45	28	62	57	32	79	30	11	53
1997/1999	39	24	55	47	25	68	30	12	52
1998/2000	41	25	59	50	26	74	31	11	54
1999/2001	45	28	62	51	27	74	37	15	61
2000/2002	55	37	70	54	28	78	56	32	75
2001/2003	51	35	66	47	23	71	54	33	72
Alberta									
1992/1994	51	40	62	50	35	64	54	35	70
1993/1995	49	37	61	52	37	67	44	26	62
1994/1996	47	35	60	49	33	64	44	23	64
1995/1997	48	35	61	39	23	55	60	39	78
1996/1998	51	39	62	46	30	61	56	39	72
1997/1999	53	42	63	51	36	65	55	39	69
1998/2000	53	43	62	57	43	70	48	34	62
1999/2001	55	45	64	57	43	69	53	38	66
2000/2002	53	43	62	54	42	66	51	36	64
2001/2003	52	42	61	54	41	66	49	34	63
British Columbia									
1992/1994	42	31	54	48	31	64	36	21	52
1993/1995	37	27	47	34	21	50	39	25	54
1994/1996	40	30	50	30	17	44	49	34	63
1995/1997	40	29	50	36	22	51	44	29	58
1996/1998	40	29	51	37	23	51	45	27	62
1997/1999	36	26	47	37	24	51	33	18	50
1998/2000	38	28	48	40	28	53	34	20	50
1999/2001	40	31	49	46	33	58	32	20	46
2000/2002	43	34	52	52	39	64	32	21	44
2001/2003	40	32	49	48	35	61	33	22	44

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).