

Table 14-3

Colon excluding rectum — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Colon excluding rectum (ICD-O-3: C18.0-C18.9, C26.0) ³									
Newfoundland and Labrador									
1992/1994	62 ^E	57 ^E	67 ^E	66 ^E	59 ^E	72 ^E	59 ^E	52 ^E	65 ^E
1993/1995	62 ^E	57 ^E	66 ^E	66 ^E	59 ^E	73 ^E	57 ^E	50 ^E	63 ^E
1994/1996	62 ^E	58 ^E	67 ^E	66 ^E	60 ^E	73 ^E	58 ^E	51 ^E	64 ^E
1995/1997	64 ^E	60 ^E	69 ^E	65 ^E	58 ^E	71 ^E	64 ^E	57 ^E	70 ^E
1996/1998	66 ^E	61 ^E	70 ^E	64 ^E	58 ^E	71 ^E	68 ^E	61 ^E	74 ^E
1997/1999	64 ^E	59 ^E	68 ^E	62 ^E	56 ^E	69 ^E	65 ^E	58 ^E	71 ^E
1998/2000	57 ^E	53 ^E	62 ^E	56 ^E	50 ^E	63 ^E	59 ^E	52 ^E	65 ^E
1999/2001	57 ^E	53 ^E	62 ^E	55 ^E	49 ^E	62 ^E	59 ^E	53 ^E	66 ^E
2000/2002	60 ^E	55 ^E	64 ^E	60 ^E	54 ^E	66 ^E	60 ^E	53 ^E	66 ^E
2001/2003	66 ^E	62 ^E	70 ^E	66 ^E	60 ^E	72 ^E	66 ^E	60 ^E	72 ^E
Prince Edward Island									
1992/1994	58	49	67	61	46	75	57	45	68
1993/1995	62	52	71	67	50	83	58	46	69
1994/1996	57	48	66	54	38	70	59	47	69
1995/1997	62	53	72	61	46	76	63	51	74
1996/1998	60	51	68	57	43	71	62	50	72
1997/1999	58	49	67	64	51	77	52	41	64
1998/2000	54	45	63	67	52	80	43	31	54
1999/2001	55	45	64	68	53	83	43	32	55
2000/2002	59	50	68	65	50	80	55	43	66
2001/2003	57	48	66	61	46	76	55	44	66
Nova Scotia									
1992/1994	55	52	59	55	50	60	56	50	60
1993/1995	53	50	57	54	49	59	53	48	58
1994/1996	53	49	56	54	49	59	52	47	56
1995/1997	53	49	56	55	50	61	50	45	55
1996/1998	55	51	58	59	54	64	51	46	55
1997/1999	55	52	59	57	52	62	54	49	58
1998/2000	55	51	58	53	48	58	56	52	61
1999/2001	57	53	60	51	46	56	62	57	67
2000/2002	55	52	58	50	46	55	60	55	64
2001/2003	58	55	61	55	50	59	62	57	66
New Brunswick									
1992/1994	53	49	57	56	50	61	51	46	57
1993/1995	54	50	58	57	51	63	52	46	57
1994/1996	55	51	59	55	49	61	54	49	60
1995/1997	55	51	59	55	49	61	55	50	61
1996/1998	56	52	60	54	48	60	58	52	63
1997/1999	58	54	61	55	49	61	60	54	65
1998/2000	59	55	63	56	50	61	62	56	67
1999/2001	59	55	63	58	52	64	60	55	66
2000/2002	60	56	64	58	52	64	62	57	68
2001/2003	60	56	64	58	52	64	61	56	66
Ontario									
1992/1994	56	55	57	55	53	56	56	55	58
1993/1995	57	55	58	57	55	58	56	55	58
1994/1996	58	57	59	58	56	59	58	57	60
1995/1997	58	57	60	58	56	59	59	58	61
1996/1998	58	57	59	58	56	59	59	58	61
1997/1999	58	57	59	58	56	59	59	57	60
1998/2000	59	58	60	59	58	61	59	57	60
1999/2001	61	60	62	61	59	62	61	60	62
2000/2002	62	61	63	62	60	63	63	61	64
2001/2003	64	63	65	63	62	65	64	63	66

See footnotes at the end of the table.

Table 14-3 – continued

Colon excluding rectum — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	58	55	62	58	53	62	59	54	63
1993/1995	58	55	62	59	54	64	58	53	62
1994/1996	55	52	58	57	52	62	53	48	58
1995/1997	56	52	59	58	53	63	53	49	58
1996/1998	57	54	61	59	54	63	56	51	60
1997/1999	58	55	62	57	52	61	60	55	64
1998/2000	58	55	61	57	53	62	59	55	64
1999/2001	54	51	57	52	48	57	56	52	61
2000/2002	56	53	59	55	51	60	57	52	61
2001/2003	59	56	62	58	54	63	59	55	64
Saskatchewan									
1992/1994	55	51	59	52	47	57	59	53	64
1993/1995	55	51	58	52	47	57	57	52	63
1994/1996	55	51	58	54	48	59	56	51	61
1995/1997	58	54	62	57	52	62	58	53	64
1996/1998	59	55	62	57	52	62	60	55	65
1997/1999	60	56	63	56	51	61	63	58	68
1998/2000	58	55	62	58	53	63	59	54	64
1999/2001	59	55	62	60	54	65	58	53	63
2000/2002	59	55	62	60	55	65	57	52	62
2001/2003	58	55	62	60	55	65	57	52	62
Alberta									
1992/1994	55	52	58	53	49	57	57	53	61
1993/1995	56	53	59	53	50	57	59	55	62
1994/1996	56	53	59	54	51	58	58	54	62
1995/1997	55	52	57	54	50	57	56	53	60
1996/1998	54	52	57	54	50	57	55	51	58
1997/1999	56	53	58	56	53	60	55	51	58
1998/2000	58	55	60	59	55	62	56	53	60
1999/2001	58	56	60	59	56	62	57	53	60
2000/2002	58	56	61	59	56	62	58	55	61
2001/2003	57	55	59	59	55	62	56	52	59
British Columbia									
1992/1994	56	54	59	56	53	59	57	54	60
1993/1995	55	53	57	54	51	57	56	53	59
1994/1996	56	54	58	55	52	58	57	54	60
1995/1997	55	53	57	53	50	56	58	55	60
1996/1998	56	54	58	55	52	58	57	55	60
1997/1999	56	55	58	55	53	58	58	55	60
1998/2000	57	55	59	57	54	60	57	54	60
1999/2001	59	57	61	59	56	62	59	56	61
2000/2002	59	57	61	60	57	62	58	56	61
2001/2003	60	58	62	60	57	63	60	57	62

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).