

Table 20-3

Other digestive system — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Other digestive system (ICD-O-3: C22.1, C24.0-C24.9, C26.8-C26.9, C48.0, C48.1-C48.2, C48.8) 3									
Newfoundland and Labrador									
1992/1994	35 E	16 E	57 E	35 E	11 E	65 E	F	F	F
1993/1995	28 E	10 E	51 E	35 E	8 E	69 E	22 E	3 E	52 E
1994/1996	8 E	1 E	23 E	16 E	3 E	42 E	0 E	..	..
1995/1997	8 E	1 E	22 E	9 E	1 E	33 E	7 E	0 E	28 E
1996/1998	23 E	9 E	42 E	18 E	3 E	45 E	27 E	9 E	53 E
1997/1999	30 E	13 E	50 E	10 E	1 E	38 E	45 E	18 E	71 E
1998/2000	40 E	21 E	60 E	27 E	6 E	55 E	51 E	23 E	77 E
1999/2001	36 E	19 E	55 E	28 E	9 E	53 E	45 E	18 E	72 E
2000/2002	36 E	20 E	54 E	35 E	15 E	59 E	37 E	15 E	62 E
2001/2003	25 E	12 E	40 E	26 E	10 E	48 E	23 E	7 E	46 E
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	12	1	43	F	F	F	F	F	F
1995/1997	12	1	43	F	F	F	F	F	F
1996/1998	15	3	40	F	F	F	F	F	F
1997/1999	10	1	36	F	F	F	F	F	F
1998/2000	7	0	28	F	F	F	0	..	..
1999/2001	0	..	..	F	F	F	0	..	..
2000/2002	0	..	..	F	F	F	0	..	..
2001/2003	12	2	32	F	F	F	19	3	46
Nova Scotia									
1992/1994	18	10	29	11	3	26	24	12	38
1993/1995	12	6	21	10	3	24	14	6	27
1994/1996	10	4	18	3	0	15	15	6	27
1995/1997	10	4	20	4	0	17	16	6	30
1996/1998	12	6	22	4	0	15	20	9	34
1997/1999	13	7	22	13	4	27	14	6	26
1998/2000	11	5	19	16	6	31	8	3	17
1999/2001	11	5	19	15	5	29	8	3	18
2000/2002	13	7	21	10	3	24	15	7	26
2001/2003	16	10	24	21	10	36	13	6	23
New Brunswick									
1992/1994	14	6	24	17	6	33	11	3	25
1993/1995	14	6	24	10	3	25	17	6	33
1994/1996	12	5	23	8	1	23	16	5	32
1995/1997	12	5	21	9	2	21	15	5	32
1996/1998	13	6	24	10	3	24	19	6	38
1997/1999	12	5	22	12	4	25	12	3	29
1998/2000	15	7	27	21	8	39	9	1	24
1999/2001	11	5	21	17	5	35	6	1	19
2000/2002	8	3	16	13	3	30	5	1	15
2001/2003	6	2	14	5	0	20	8	2	17
Ontario									
1992/1994	20	17	22	18	15	22	21	18	25
1993/1995	22	19	24	21	18	25	22	19	26
1994/1996	21	19	24	21	17	25	22	18	26
1995/1997	20	18	23	20	16	23	20	17	24
1996/1998	18	15	20	17	14	20	19	16	22
1997/1999	18	16	20	18	15	21	18	15	21
1998/2000	18	16	20	19	16	23	17	14	20
1999/2001	21	18	23	21	18	25	20	16	23
2000/2002	21	18	23	21	18	24	20	17	24
2001/2003	22	20	25	21	18	25	24	20	27

See footnotes at the end of the table.

Table 20-3 – continued

Other digestive system — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	11	6	19	11	4	22	12	4	24
1993/1995	11	6	18	13	6	24	9	3	20
1994/1996	13	7	20	15	7	26	11	4	22
1995/1997	10	6	17	13	6	24	8	2	17
1996/1998	16	10	24	17	9	29	15	7	26
1997/1999	17	11	24	17	8	29	17	9	27
1998/2000	17	11	24	16	8	26	18	10	29
1999/2001	14	9	21	11	5	20	18	9	29
2000/2002	16	10	23	10	4	20	21	12	33
2001/2003	18	11	26	11	5	22	23	13	35
Saskatchewan									
1992/1994	16	9	25	15	6	29	16	7	29
1993/1995	19	12	28	20	9	33	19	9	31
1994/1996	13	7	22	11	4	24	15	6	27
1995/1997	17	10	27	14	5	27	21	10	35
1996/1998	17	10	27	9	2	21	24	13	39
1997/1999	18	10	27	10	3	22	25	13	39
1998/2000	18	11	26	16	7	30	18	9	30
1999/2001	20	13	29	20	9	34	20	11	32
2000/2002	18	11	26	19	9	32	17	9	27
2001/2003	19	13	27	14	6	26	23	14	34
Alberta									
1992/1994	18	13	24	21	13	30	15	9	23
1993/1995	18	13	24	23	15	33	14	8	22
1994/1996	19	14	24	22	14	31	16	10	23
1995/1997	16	12	21	16	10	24	16	10	23
1996/1998	14	10	19	10	5	17	17	11	23
1997/1999	13	9	17	9	4	17	14	10	20
1998/2000	13	10	18	9	4	16	16	11	21
1999/2001	14	10	18	11	6	17	16	11	21
2000/2002	14	11	18	12	7	18	16	11	21
2001/2003	15	11	19	16	10	23	14	10	20
British Columbia									
1992/1994	15	10	20	17	11	25	13	7	20
1993/1995	15	10	20	16	10	24	14	8	21
1994/1996	15	11	20	17	11	24	14	8	20
1995/1997	14	11	19	18	12	25	11	7	17
1996/1998	17	13	21	17	11	24	17	12	23
1997/1999	17	13	21	14	9	20	19	14	25
1998/2000	15	11	19	11	7	16	18	13	24
1999/2001	11	8	14	10	6	15	12	8	16
2000/2002	10	8	14	11	7	17	10	7	14
2001/2003	13	10	17	16	11	23	11	8	16

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).