

Table 23-3

Other respiratory system — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Other respiratory system (ICD-O-3: C30.0-C30.1, C31.0-C31.9, C33.9, C38.1-C38.3, C38.4, C38.8, C39.0, C39.8, C39.9) ³									
Newfoundland and Labrador									
1992/1994	44 ^E	18 ^E	70 ^E	F	F	F	F	F	F
1993/1995	46 ^E	21 ^E	69 ^E	F	F	F	F	F	F
1994/1996	52 ^E	23 ^E	78 ^E	F	F	F	F	F	F
1995/1997	47 ^E	14 ^E	80 ^E	F	F	F	F	F	F
1996/1998	44 ^E	14 ^E	78 ^E	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	36 ^E	9 ^E	70 ^E	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	44 ^E	13 ^E	73 ^E	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	..	..	..
Nova Scotia									
1992/1994	52	28	73	39	12	67	67	28	92
1993/1995	39	17	62	41	15	69	F	F	F
1994/1996	50	27	72	38	16	62	F	F	F
1995/1997	50	27	72	42	17	68	F	F	F
1996/1998	71	41	94	53	19	85	F	F	F
1997/1999	68	36	92	54	16	90	F	F	F
1998/2000	66	30	93	F	F	F	F	F	F
1999/2001	38	17	60	11	1	39	66	28	91
2000/2002	33	14	53	10	1	37	55	23	80
2001/2003	27	12	46	19	5	43	39	12	68
New Brunswick									
1992/1994	65	30	91	74	31	103	F	F	F
1993/1995	47	19	72	66	28	91	F	F	F
1994/1996	38	14	63	F	F	F	F	F	F
1995/1997	27	8	50	F	F	F	F	F	F
1996/1998	41	17	65	F	F	F	39	12	67
1997/1999	34	10	62	F	F	F	33	8	64
1998/2000	49	18	78	F	F	F	F	F	F
1999/2001	26	8	50	36	9	70	F	F	F
2000/2002	37	15	61	44	14	74	F	F	F
2001/2003	41	20	63	45	18	70	F	F	F
Ontario									
1992/1994	40	35	45	38	32	44	45	36	54
1993/1995	40	35	45	41	35	48	37	29	46
1994/1996	41	36	46	41	35	48	40	31	49
1995/1997	41	36	46	42	35	48	40	32	48
1996/1998	40	35	45	39	32	45	41	33	49
1997/1999	42	36	47	43	36	50	40	32	48
1998/2000	39	34	45	37	30	44	43	34	52
1999/2001	41	35	47	40	32	47	43	33	52
2000/2002	39	33	44	34	28	41	46	37	55
2001/2003	39	33	44	37	30	44	42	33	55

See footnotes at the end of the table.

Table 23-3 – continued

Other respiratory system — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	22	10	38	18	6	36	31	7	61
1993/1995	22	10	39	10	2	28	42	15	70
1994/1996	32	17	48	17	5	36	61	28	86
1995/1997	41	24	57	22	7	44	62	34	83
1996/1998	43	26	60	36	16	57	55	25	79
1997/1999	49	30	67	50	24	73	47	20	73
1998/2000	59	37	77	66	38	86	45	14	76
1999/2001	55	32	76	66	35	89	F	F	F
2000/2002	56	34	77	59	29	85	53	20	83
2001/2003	49	26	72	46	19	74	F	F	F
Saskatchewan									
1992/1994	39	17	63	44	18	68	F	F	F
1993/1995	44	22	65	49	22	73	F	F	F
1994/1996	50	30	69	56	31	78	F	F	F
1995/1997	40	22	58	41	20	63	F	F	F
1996/1998	37	18	56	45	20	69	22	3	52
1997/1999	30	12	52	24	6	51	F	F	F
1998/2000	35	16	58	35	11	65	36	8	69
1999/2001	51	27	74	47	17	77	56	21	87
2000/2002	44	23	64	43	16	71	45	16	73
2001/2003	62	36	82	46	14	78	75	38	97
Alberta									
1992/1994	44	31	57	54	34	72	33	17	51
1993/1995	36	23	50	43	25	62	27	11	47
1994/1996	36	23	49	44	27	61	23	8	43
1995/1997	40	28	53	47	31	62	27	10	49
1996/1998	42	30	53	47	33	61	26	10	47
1997/1999	43	31	55	43	29	57	42	20	64
1998/2000	36	24	47	32	20	46	44	21	65
1999/2001	34	22	46	25	13	39	56	29	78
2000/2002	46	32	59	38	23	53	68	36	90
2001/2003	54	40	66	46	31	61	73	44	91
British Columbia									
1992/1994	55	43	66	48	32	64	62	44	77
1993/1995	60	48	70	57	43	70	64	45	80
1994/1996	49	38	60	45	31	59	57	38	73
1995/1997	46	35	56	42	29	55	52	35	67
1996/1998	37	27	48	24	13	38	51	35	66
1997/1999	45	34	55	38	24	53	52	36	66
1998/2000	41	30	52	40	25	55	42	27	56
1999/2001	49	38	60	58	40	73	42	27	56
2000/2002	50	39	60	57	41	72	42	28	57
2001/2003	54	43	65	59	43	73	49	33	64

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).