

Table 25-3

Soft tissue (including heart) — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Soft tissue (including heart) (ICD-O-3: C38.0, C47.0-C47.9, C49.0-C49.9) ³									
Newfoundland and Labrador									
1992/1994	55 ^E	32 ^E	76 ^E	56 ^E	26 ^E	83 ^E	53 ^E	19 ^E	80 ^E
1993/1995	61 ^E	38 ^E	79 ^E	52 ^E	22 ^E	79 ^E	69 ^E	35 ^E	89 ^E
1994/1996	57 ^E	36 ^E	75 ^E	53 ^E	26 ^E	76 ^E	62 ^E	29 ^E	86 ^E
1995/1997	63 ^E	44 ^E	78 ^E	61 ^E	35 ^E	82 ^E	65 ^E	36 ^E	85 ^E
1996/1998	58 ^E	40 ^E	74 ^E	59 ^E	33 ^E	80 ^E	57 ^E	31 ^E	79 ^E
1997/1999	64 ^E	46 ^E	78 ^E	68 ^E	40 ^E	88 ^E	60 ^E	36 ^E	79 ^E
1998/2000	52 ^E	34 ^E	68 ^E	49 ^E	22 ^E	73 ^E	54 ^E	30 ^E	75 ^E
1999/2001	51 ^E	35 ^E	65 ^E	42 ^E	22 ^E	61 ^E	61 ^E	37 ^E	80 ^E
2000/2002	50 ^E	33 ^E	65 ^E	30 ^E	12 ^E	51 ^E	70 ^E	44 ^E	88 ^E
2001/2003	60 ^E	43 ^E	75 ^E	35 ^E	14 ^E	58 ^E	80 ^E	55 ^E	94 ^E
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	21	3	49	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Nova Scotia									
1992/1994	63	48	76	66	45	83	59	37	77
1993/1995	63	48	76	68	47	85	58	36	76
1994/1996	55	40	68	58	37	76	51	30	70
1995/1997	47	33	61	46	27	64	50	28	69
1996/1998	51	36	65	51	31	69	52	29	72
1997/1999	60	44	74	61	39	79	59	34	79
1998/2000	64	48	78	66	45	83	62	37	81
1999/2001	60	45	73	59	39	76	61	37	79
2000/2002	56	41	70	61	42	78	48	26	69
2001/2003	47	33	61	50	31	68	44	23	64
New Brunswick									
1992/1994	55	39	69	62	37	84	49	29	67
1993/1995	52	34	70	64	36	86	41	18	64
1994/1996	60	41	76	68	39	89	52	28	73
1995/1997	59	42	73	55	28	77	61	39	79
1996/1998	62	48	74	61	40	79	63	44	79
1997/1999	62	48	74	61	39	80	62	44	77
1998/2000	60	45	73	61	39	80	59	39	76
1999/2001	59	44	72	52	30	72	65	44	81
2000/2002	67	52	81	62	41	80	75	51	91
2001/2003	66	51	80	66	46	84	66	43	85
Ontario									
1992/1994	63	59	67	62	57	68	63	57	68
1993/1995	63	59	67	64	59	69	62	56	67
1994/1996	63	59	67	65	59	70	62	56	67
1995/1997	62	58	66	61	56	66	64	58	69
1996/1998	61	57	64	60	54	65	62	56	67
1997/1999	58	54	62	57	51	62	60	54	65
1998/2000	60	56	63	60	55	65	59	54	64
1999/2001	62	58	65	62	57	67	62	56	66
2000/2002	63	59	66	63	58	68	62	57	67
2001/2003	65	62	69	66	61	71	64	59	69

See footnotes at the end of the table.

Table 25-3 – continued

Soft tissue (including heart) — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	57	43	69	65	47	81	45	25	64
1993/1995	61	47	73	69	51	84	49	30	67
1994/1996	57	44	69	68	50	83	43	25	61
1995/1997	60	46	72	66	48	82	49	29	67
1996/1998	59	45	72	66	46	83	50	31	68
1997/1999	66	51	79	69	48	87	63	41	80
1998/2000	55	42	68	51	34	67	64	42	80
1999/2001	58	45	69	55	39	71	62	42	78
2000/2002	55	44	65	56	40	70	54	38	69
2001/2003	58	46	69	58	41	73	58	42	73
Saskatchewan									
1992/1994	67	49	81	81	57	97	49	25	71
1993/1995	65	48	79	88	63	103	41	21	60
1994/1996	60	44	73	77	56	92	34	15	55
1995/1997	56	41	69	64	45	80	42	22	63
1996/1998	53	39	66	54	36	70	51	29	71
1997/1999	62	48	75	68	50	82	52	28	73
1998/2000	62	49	74	68	52	82	51	31	70
1999/2001	69	57	80	74	57	86	63	43	79
2000/2002	64	51	75	57	39	73	71	53	85
2001/2003	66	53	78	62	42	80	70	51	85
Alberta									
1992/1994	66	55	74	57	42	70	75	60	85
1993/1995	67	57	76	62	47	74	73	58	84
1994/1996	67	57	76	57	42	71	78	63	88
1995/1997	59	49	68	58	44	70	60	46	73
1996/1998	56	47	65	54	41	65	59	45	72
1997/1999	59	50	67	57	45	67	62	48	74
1998/2000	71	62	78	67	55	77	78	63	88
1999/2001	70	61	77	68	56	78	71	59	81
2000/2002	67	58	74	68	56	78	65	53	76
2001/2003	59	50	66	56	45	67	61	49	72
British Columbia									
1992/1994	64	55	72	60	48	70	71	57	81
1993/1995	66	58	73	63	52	73	69	57	79
1994/1996	65	57	72	63	53	73	66	55	75
1995/1997	63	57	69	61	51	69	66	56	75
1996/1998	65	59	71	61	52	69	71	61	79
1997/1999	65	58	71	60	51	68	71	61	79
1998/2000	70	63	76	66	56	74	74	64	83
1999/2001	72	65	78	72	62	81	72	60	81
2000/2002	73	66	79	77	67	85	69	58	78
2001/2003	74	67	80	75	66	82	73	61	82

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).