

Table 34-1

Testis — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	1 year of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Testis (ICD-O-3: C62.0-C62.9) ²									
1992									
Total, 15 to 99 years	.	.	.	96	94	98	.	.	.
15 to 44 years	.	.	.	96	93	98	.	.	.
45 to 54 years	.	.	.	102	..	..	.	.	.
55 to 64 years	.	.	.	99	65	105	.	.	.
65 to 74 years	.	.	.	83	49	103	.	.	.
75 to 99 years	.	.	.	F	F	F	.	.	.
1993									
Total, 15 to 99 years	.	.	.	94	91	96	.	.	.
15 to 44 years	.	.	.	96	93	97	.	.	.
45 to 54 years	.	.	.	91	76	97	.	.	.
55 to 64 years	.	.	.	89	61	101	.	.	.
65 to 74 years	.	.	.	69	32	95	.	.	.
75 to 99 years	.	.	.	F	F	F	.	.	.
1994									
Total, 15 to 99 years	.	.	.	95	93	97	.	.	.
15 to 44 years	.	.	.	96	94	98	.	.	.
45 to 54 years	.	.	.	93	83	98	.	.	.
55 to 64 years	.	.	.	91	67	101	.	.	.
65 to 74 years	.	.	.	85	52	104	.	.	.
75 to 99 years	.	.	.	..	..	..	.	.	.
1995									
Total, 15 to 99 years	.	.	.	95	92	97	.	.	.
15 to 44 years	.	.	.	96	93	97	.	.	.
45 to 54 years	.	.	.	91	78	97	.	.	.
55 to 64 years	.	.	.	100	69	105	.	.	.
65 to 74 years	.	.	.	96	49	113	.	.	.
75 to 99 years	.	.	.	F	F	F	.	.	.
1996									
Total, 15 to 99 years	.	.	.	95	93	97	.	.	.
15 to 44 years	.	.	.	96	94	98	.	.	.
45 to 54 years	.	.	.	94	84	98	.	.	.
55 to 64 years	.	.	.	106	..	..	.	.	.
65 to 74 years	.	.	.	F	F	F	.	.	.
75 to 99 years	.	.	.	F	F	F	.	.	.
1997									
Total, 15 to 99 years	.	.	.	94	92	96	.	.	.
15 to 44 years	.	.	.	95	93	97	.	.	.
45 to 54 years	.	.	.	96	87	100	.	.	.
55 to 64 years	.	.	.	71	40	90	.	.	.
65 to 74 years	.	.	.	F	F	F	.	.	.
75 to 99 years	.	.	.	72	26	108	.	.	.
1998									
Total, 15 to 99 years	.	.	.	97	95	98	.	.	.
15 to 44 years	.	.	.	98	96	99	.	.	.
45 to 54 years	.	.	.	95	84	99	.	.	.
55 to 64 years	.	.	.	81	44	98	.	.	.
65 to 74 years	.	.	.	92	47	109	.	.	.
75 to 99 years	.	.	.	F	F	F	.	.	.
1999									
Total, 15 to 99 years	.	.	.	96	93	97	.	.	.
15 to 44 years	.	.	.	96	94	98	.	.	.
45 to 54 years	.	.	.	99	91	101	.	.	.
55 to 64 years	.	.	.	96	72	103	.	.	.
65 to 74 years	.	.	.	74	34	98	.	.	.
75 to 99 years	.	.	.	F	F	F	.	.	.

See footnotes at the end of the table.

Testis — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

See footnotes at the end of the table.

Table 34-1 – continued

Testis — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Testis (ICD-O-3: C62.0-C62.9) 2									
1992/1994									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	95	89	98	.	.	.
55 to 64 years	.	.	.	93	80	99	.	.	.
65 to 74 years	.	.	.	80	61	94	.	.	.
75 to 99 years	.	.	.	72	22	124	.	.	.
1993/1995									
Total, 15 to 99 years	.	.	.	95	93	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	92	86	96	.	.	.
55 to 64 years	.	.	.	93	81	100	.	.	.
65 to 74 years	.	.	.	83	62	97	.	.	.
75 to 99 years	.	.	.	46	11	94	.	.	.
1994/1996									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	93	87	96	.	.	.
55 to 64 years	.	.	.	98	86	103	.	.	.
65 to 74 years	.	.	.	83	62	97	.	.	.
75 to 99 years	.	.	.	61	22	100	.	.	.
1995/1997									
Total, 15 to 99 years	.	.	.	95	93	96	.	.	.
15 to 44 years	.	.	.	96	94	97	.	.	.
45 to 54 years	.	.	.	94	89	97	.	.	.
55 to 64 years	.	.	.	92	78	100	.	.	.
65 to 74 years	.	.	.	80	55	96	.	.	.
75 to 99 years	.	.	.	66	35	94	.	.	.
1996/1998									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	95	90	98	.	.	.
55 to 64 years	.	.	.	86	70	96	.	.	.
65 to 74 years	.	.	.	78	55	94	.	.	.
75 to 99 years	.	.	.	74	40	103	.	.	.
1997/1999									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	97	93	99	.	.	.
55 to 64 years	.	.	.	85	70	94	.	.	.
65 to 74 years	.	.	.	81	58	96	.	.	.
75 to 99 years	.	.	.	59	28	89	.	.	.
1998/2000									
Total, 15 to 99 years	.	.	.	96	95	97	.	.	.
15 to 44 years	.	.	.	97	96	98	.	.	.
45 to 54 years	.	.	.	97	93	100	.	.	.
55 to 64 years	.	.	.	93	81	100	.	.	.
65 to 74 years	.	.	.	77	55	92	.	.	.
75 to 99 years	.	.	.	56	23	91	.	.	.
1999/2001									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	95	91	98	.	.	.
55 to 64 years	.	.	.	95	83	100	.	.	.
65 to 74 years	.	.	.	75	53	91	.	.	.
75 to 99 years	.	.	.	67	39	94	.	.	.

See footnotes at the end of the table.

Table 34-1 – continued

Testis — Five-year relative survival ratios, October 2011 CCR file, by age group and sex, 1 and 3 years of cases, Canada (excluding Quebec)¹

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
2000/2002									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	93	89	96	.	.	.
55 to 64 years	.	.	.	94	83	100	.	.	.
65 to 74 years	.	.	.	78	54	94	.	.	.
75 to 99 years	.	.	.	70	39	98	.	.	.
2001/2003									
Total, 15 to 99 years	.	.	.	95	94	96	.	.	.
15 to 44 years	.	.	.	96	95	97	.	.	.
45 to 54 years	.	.	.	94	90	97	.	.	.
55 to 64 years	.	.	.	89	77	97	.	.	.
65 to 74 years	.	.	.	99	76	109	.	.	.
75 to 99 years	.	.	.	74	43	103	.	.	.

1. Data from Quebec have been excluded, in part, because the method of ascertaining the date of cancer diagnosis differs from the method used by other registries and because of issues in correctly ascertaining the vital status of cases.

2. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM tables 103-1565 and 103-1566).