

Table 39-3

Ureter — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Ureter (ICD-O-3: C66.9) ³									
Newfoundland and Labrador									
1992/1994	65 ^E	30 ^E	91 ^E	67 ^E	28 ^E	92 ^E	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Nova Scotia									
1992/1994	53	30	76	64	35	88	F	F	F
1993/1995	67	41	88	71	42	93	F	F	F
1994/1996	59	33	82	62	32	87	F	F	F
1995/1997	79	49	99	80	43	104	F	F	F
1996/1998	84	51	103	82	41	106	F	F	F
1997/1999	78	49	97	79	44	102	F	F	F
1998/2000	54	30	76	49	20	77	F	F	F
1999/2001	50	28	72	53	22	83	46	17	76
2000/2002	56	33	77	59	29	87	50	18	80
2001/2003	61	37	83	68	37	93	50	15	83
New Brunswick									
1992/1994	92	47	109	F	F	F	F	F	F
1993/1995	85	46	103	F	F	F	F	F	F
1994/1996	65	30	90	F	F	F	F	F	F
1995/1997	45	14	80	55	17	92	F	F	F
1996/1998	42	13	75	F	F	F	F	F	F
1997/1999	38	9	75	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	50	20	78	F	F	F	F	F	F
Ontario									
1992/1994	56	45	67	60	46	72	49	29	68
1993/1995	52	41	63	56	42	69	45	26	64
1994/1996	55	44	65	61	47	73	42	25	61
1995/1997	61	49	71	73	58	86	39	23	55
1996/1998	63	52	73	66	53	79	57	39	72
1997/1999	58	47	68	59	46	72	55	37	71
1998/2000	53	43	63	52	39	65	54	37	70
1999/2001	54	44	64	54	40	67	55	39	70
2000/2002	55	44	64	52	39	65	57	42	71
2001/2003	55	45	65	50	36	63	62	46	77

See footnotes at the end of the table.

Table 39-3 – continued

Ureter — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	57	28	82	47	14	79	F	F	F
1993/1995	69	42	90	63	26	90	75	35	103
1994/1996	69	43	90	66	28	94	72	36	98
1995/1997	51	25	77	48	15	82	54	17	91
1996/1998	35	13	62	36	9	72	F	F	F
1997/1999	57	32	80	53	24	81	F	F	F
1998/2000	71	46	90	62	32	88	F	F	F
1999/2001	71	47	90	61	33	85	F	F	F
2000/2002	69	41	90	63	31	89	F	F	F
2001/2003	62	32	86	54	22	83	F	F	F
Saskatchewan									
1992/1994	53	26	77	63	23	94	F	F	F
1993/1995	75	45	96	78	36	103	F	F	F
1994/1996	70	39	93	F	F	F	F	F	F
1995/1997	65	35	88	73	34	100	F	F	F
1996/1998	53	26	78	72	33	98	F	F	F
1997/1999	61	33	85	74	39	98	F	F	F
1998/2000	57	26	84	66	28	95	F	F	F
1999/2001	66	33	91	69	29	95	F	F	F
2000/2002	82	47	101	F	F	F	F	F	F
2001/2003	83	47	102	F	F	F	F	F	F
Alberta									
1992/1994	60	35	81	68	36	92	F	F	F
1993/1995	63	38	83	65	36	88	F	F	F
1994/1996	74	49	92	74	43	95	F	F	F
1995/1997	67	44	85	68	38	90	64	30	91
1996/1998	72	51	89	82	51	102	62	32	85
1997/1999	53	33	72	66	37	91	32	10	59
1998/2000	54	32	76	68	37	94	31	7	61
1999/2001	45	25	64	51	24	76	36	11	65
2000/2002	48	29	67	44	20	70	52	24	78
2001/2003	48	31	65	46	26	68	50	22	76
British Columbia									
1992/1994	54	37	70	62	38	82	42	19	66
1993/1995	42	27	57	45	26	64	37	17	59
1994/1996	34	20	48	40	22	58	23	7	44
1995/1997	37	23	51	38	21	58	34	15	56
1996/1998	43	29	57	44	26	62	42	21	64
1997/1999	42	28	55	35	19	54	50	28	70
1998/2000	39	26	54	38	21	58	40	20	60
1999/2001	45	29	60	39	19	61	51	29	72
2000/2002	58	43	71	56	38	73	62	35	82
2001/2003	55	41	67	48	33	63	74	46	93

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).