

Table 45-3

Other endocrine — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Other endocrine (ICD-O-3: C37.9, C74.0-C74.9, C75.0-C75.9) ³									
Newfoundland and Labrador									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	..	..	..	F	F	F
Nova Scotia									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	46	17	73	F	F	F	F	F	F
1995/1997	59	25	84	F	F	F	F	F	F
1996/1998	64	27	88	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	42	13	71	F	F	F	F	F	F
1999/2001	52	24	75	F	F	F	F	F	F
2000/2002	65	36	85	F	F	F	84	43	99
2001/2003	72	39	91	F	F	F	F	F	F
New Brunswick									
1992/1994	57	24	81	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	58	27	81	F	F	F	F	F	F
1997/1999	53	22	78	F	F	F	F	F	F
1998/2000	70	35	90	F	F	F	F	F	F
1999/2001	86	47	100	F	F	F	F	F	F
2000/2002	72	39	91	F	F	F	F	F	F
2001/2003	48	18	75	F	F	F	F	F	F
Ontario									
1992/1994	55	48	62	46	36	55	67	56	76
1993/1995	63	56	70	55	45	64	72	62	80
1994/1996	69	62	76	65	55	74	74	63	83
1995/1997	65	58	72	60	50	69	72	61	81
1996/1998	64	57	71	64	53	73	65	54	75
1997/1999	57	50	64	57	47	66	58	48	67
1998/2000	61	55	68	59	49	68	63	54	72
1999/2001	61	54	68	54	44	64	68	58	76
2000/2002	64	57	71	50	40	59	76	67	83
2001/2003	65	58	71	55	45	65	72	63	80

See footnotes at the end of the table.

Table 45-3 – continued

Other endocrine — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	39	17	61	57	21	86	22	3	51
1993/1995	41	19	64	F	F	F	34	8	65
1994/1996	44	18	69	F	F	F	F	F	F
1995/1997	51	21	77	F	F	F	F	F	F
1996/1998	57	26	80	F	F	F	F	F	F
1997/1999	48	20	73	F	F	F	F	F	F
1998/2000	53	22	78	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Saskatchewan									
1992/1994	54	28	76	F	F	F	F	F	F
1993/1995	58	26	81	F	F	F	F	F	F
1994/1996	53	26	75	F	F	F	53	19	80
1995/1997	65	32	86	F	F	F	F	F	F
1996/1998	68	40	87	F	F	F	74	35	95
1997/1999	68	36	88	F	F	F	F	F	F
1998/2000	69	37	89	F	F	F	F	F	F
1999/2001	66	33	87	F	F	F	F	F	F
2000/2002	46	21	69	F	F	F	F	F	F
2001/2003	45	19	70	F	F	F	62	26	85
Alberta									
1992/1994	41	22	60	43	16	68	40	14	65
1993/1995	42	22	61	53	22	78	32	10	57
1994/1996	43	18	67	F	F	F	F	F	F
1995/1997	53	30	72	72	37	93	34	11	62
1996/1998	50	32	67	66	37	86	36	15	59
1997/1999	55	39	68	68	46	84	40	20	59
1998/2000	55	39	69	69	45	85	41	21	60
1999/2001	57	41	71	69	45	85	45	24	64
2000/2002	71	55	83	80	55	94	64	42	80
2001/2003	69	54	81	74	49	89	65	44	81
British Columbia									
1992/1994	46	28	63	58	29	81	37	17	59
1993/1995	49	33	64	60	37	79	35	14	57
1994/1996	55	40	68	56	35	73	54	32	72
1995/1997	62	48	74	66	46	81	58	36	74
1996/1998	64	49	76	72	50	88	56	37	73
1997/1999	61	46	73	69	46	85	54	34	71
1998/2000	62	47	75	67	43	85	58	37	76
1999/2001	76	61	87	75	52	90	77	55	91
2000/2002	68	54	80	78	55	93	60	41	75
2001/2003	62	48	74	66	42	84	59	40	75

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).