

Table 1-3

All primary sites of cancer combined — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2,3}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
All primary sites of cancer combined									
Newfoundland and Labrador									
1992/1994	57 ^E	56 ^E	59 ^E	54 ^E	52 ^E	56 ^E	61 ^E	59 ^E	63 ^E
1993/1995	58 ^E	56 ^E	59 ^E	55 ^E	53 ^E	57 ^E	61 ^E	59 ^E	63 ^E
1994/1996	59 ^E	57 ^E	60 ^E	56 ^E	53 ^E	58 ^E	62 ^E	60 ^E	65 ^E
1995/1997	60 ^E	58 ^E	61 ^E	56 ^E	54 ^E	58 ^E	64 ^E	62 ^E	66 ^E
1996/1998	62 ^E	60 ^E	63 ^E	59 ^E	57 ^E	61 ^E	64 ^E	62 ^E	67 ^E
1997/1999	61 ^E	60 ^E	63 ^E	59 ^E	56 ^E	61 ^E	65 ^E	62 ^E	67 ^E
1998/2000	61 ^E	60 ^E	63 ^E	60 ^E	58 ^E	62 ^E	63 ^E	61 ^E	65 ^E
1999/2001	62 ^E	61 ^E	63 ^E	60 ^E	58 ^E	62 ^E	65 ^E	63 ^E	67 ^E
2000/2002	63 ^E	62 ^E	65 ^E	62 ^E	60 ^E	64 ^E	65 ^E	63 ^E	67 ^E
2001/2003	63 ^E	62 ^E	65 ^E	61 ^E	59 ^E	63 ^E	66 ^E	64 ^E	68 ^E
Prince Edward Island									
1992/1994	52	50	55	50	46	54	55	51	59
1993/1995	55	52	58	52	49	56	58	54	62
1994/1996	54	51	57	50	46	54	59	55	62
1995/1997	57	54	60	53	50	57	61	57	65
1996/1998	56	53	59	53	50	57	59	55	63
1997/1999	57	54	59	56	52	60	57	54	61
1998/2000	54	52	57	54	50	58	55	51	59
1999/2001	55	53	58	57	53	60	54	50	58
2000/2002	56	54	59	58	54	61	55	51	59
2001/2003	58	56	61	60	56	63	56	53	60
Nova Scotia									
1992/1994	54	53	55	52	51	54	56	55	58
1993/1995	53	52	54	51	50	53	56	55	58
1994/1996	53	52	54	51	50	53	55	54	57
1995/1997	54	53	55	52	50	53	56	54	57
1996/1998	55	54	56	54	53	56	56	55	58
1997/1999	57	56	58	55	54	56	58	57	60
1998/2000	57	56	58	55	54	56	59	58	61
1999/2001	57	56	58	56	55	58	58	57	60
2000/2002	57	56	58	57	55	58	58	57	59
2001/2003	58	57	59	57	56	59	58	57	59
New Brunswick									
1992/1994	55	53	56	54	52	55	55	54	57
1993/1995	55	54	56	54	52	55	57	56	59
1994/1996	55	54	56	53	51	55	58	56	60
1995/1997	55	54	56	53	51	55	58	56	60
1996/1998	57	56	58	54	53	56	60	58	61
1997/1999	57	56	59	55	53	56	60	59	62
1998/2000	58	57	59	55	54	57	61	59	62
1999/2001	58	57	59	57	55	58	60	58	61
2000/2002	58	57	59	57	55	58	59	57	60
2001/2003	58	57	59	57	56	59	60	58	61
Ontario									
1992/1994	56	55	56	53	52	53	59	58	59
1993/1995	56	56	56	53	53	54	59	58	59
1994/1996	57	56	57	54	54	54	60	59	60
1995/1997	57	57	58	55	54	55	60	60	61
1996/1998	58	58	58	55	55	56	61	60	61
1997/1999	58	58	59	56	56	57	61	60	61
1998/2000	59	59	60	57	57	58	61	61	62
1999/2001	61	61	61	59	59	60	62	62	63
2000/2002	62	62	62	61	60	61	63	63	63
2001/2003	63	63	63	62	62	62	64	63	64

See footnotes at the end of the table.

Table 1-3 – continued

All primary sites of cancer combined — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2,3}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	57	56	58	57	56	58	57	55	58
1993/1995	57	56	58	57	55	58	57	55	58
1994/1996	56	55	57	54	53	56	57	56	58
1995/1997	55	54	56	53	52	54	58	56	59
1996/1998	56	55	57	54	53	55	59	57	60
1997/1999	57	56	58	55	54	56	60	59	61
1998/2000	57	56	58	55	54	56	60	59	61
1999/2001	58	57	59	56	55	58	60	59	61
2000/2002	58	57	59	58	56	59	59	58	60
2001/2003	59	58	60	58	57	60	59	58	60
Saskatchewan									
1992/1994	57	55	58	54	53	56	59	58	61
1993/1995	56	55	57	54	53	56	58	56	59
1994/1996	55	54	56	53	52	55	57	56	59
1995/1997	56	55	57	53	51	54	59	57	60
1996/1998	56	55	57	54	52	55	59	58	61
1997/1999	57	56	58	55	53	56	60	59	61
1998/2000	58	57	59	57	55	58	59	58	60
1999/2001	59	58	60	59	57	60	60	58	61
2000/2002	59	58	60	58	57	60	59	58	60
2001/2003	59	58	60	59	58	60	59	58	60
Alberta									
1992/1994	55	54	56	53	52	54	58	57	59
1993/1995	55	55	56	53	52	54	58	57	59
1994/1996	56	55	56	53	52	54	59	58	60
1995/1997	57	56	57	55	54	56	59	58	60
1996/1998	57	57	58	56	55	57	59	58	60
1997/1999	58	58	59	57	56	57	60	59	61
1998/2000	59	58	60	57	56	58	61	60	62
1999/2001	60	60	61	59	58	60	62	61	63
2000/2002	61	61	62	61	60	62	62	61	63
2001/2003	61	61	62	61	60	62	62	61	62
British Columbia									
1992/1994	58	58	59	58	57	59	59	58	60
1993/1995	58	57	58	57	56	58	59	58	60
1994/1996	57	57	58	56	55	56	59	58	60
1995/1997	57	57	58	55	55	56	59	58	60
1996/1998	58	57	58	56	56	57	59	58	60
1997/1999	58	58	59	58	57	58	59	59	60
1998/2000	59	58	59	58	57	59	60	59	60
1999/2001	60	59	60	59	59	60	60	59	61
2000/2002	60	59	60	60	59	60	60	59	61
2001/2003	60	59	60	60	59	60	60	59	61

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. Ontario does not report in situ bladder cases to the Canadian Cancer Registry.

3. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).