

Table 8-2

Oropharynx — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Oropharynx (ICD-O-3: C10.0-C10.9) ³									
1992/1994									
Canada	35	28	43	33	25	42	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	27	17	38	20	10	33	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1993/1995									
Canada	29	22	36	26	19	34	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	30	20	41	25	14	38	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1994/1996									
Canada	32	25	40	28	20	36	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	32	22	42	23	13	34	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1995/1997									
Canada	34	27	42	32	24	41	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	32	23	42	29	18	41	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1996/1998									
Canada	34	27	41	33	25	42	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	32	23	42	F	F	F	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F

See footnotes at the end of the table.

Table 8-2 – continued

Oropharynx — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
1997/1999									
Canada	28	22	36	32	23	40	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	F	F	F	F	F	F	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1998/2000									
Canada	30	23	37	30	22	38	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	F	F	F	F	F	F	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1999/2001									
Canada	30	24	37	31	24	38	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	29	20	38	31	21	42	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
2000/2002									
Canada	33	27	39	32	26	39	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	34	25	43	F	F	F	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
2001/2003									
Canada	35	29	40	35	29	42	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	..	..	..
Prince Edward Island	..	..	..	..	..	..	..	..	..
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	34	26	42	F	F	F	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1572).