

Table 8-3

Oropharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Oropharynx (ICD-O-3: C10.0-C10.9) ³									
Newfoundland and Labrador									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	..	..	..
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	..	..	..	..	..	..	..	..	..
Nova Scotia									
1992/1994	37	14	64	50	15	84	F	F	F
1993/1995	20	3	51	F	F	F	F	F	F
1994/1996	25	4	58	F	F	F	F	F	F
1995/1997	23	4	55	F	F	F	F	F	F
1996/1998	16	3	40	F	F	F	F	F	F
1997/1999	16	3	40	F	F	F	F	F	F
1998/2000	25	6	51	F	F	F	F	F	F
1999/2001	15	3	40	19	3	46	F	F	F
2000/2002	30	11	54	35	13	59	F	F	F
2001/2003	29	10	51	35	13	59	F	F	F
New Brunswick									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Ontario									
1992/1994	28	18	39	20	10	32	49	25	71
1993/1995	30	20	41	26	15	39	41	20	63
1994/1996	30	20	40	24	14	36	41	23	61
1995/1997	30	21	40	32	21	44	27	13	45
1996/1998	31	22	41	30	19	42	32	17	49
1997/1999	30	21	40	33	22	45	21	8	39
1998/2000	28	19	39	28	18	40	29	10	51
1999/2001	28	19	38	31	21	42	17	4	38
2000/2002	34	24	43	36	25	47	26	10	45
2001/2003	34	26	43	39	29	49	21	9	37

See footnotes at the end of the table.

Table 8-3 – continued

Oropharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	39	12	67	F	F	F	F	F	F
1999/2001	27	6	54	21	3	50	F	F	F
2000/2002	14	2	36	8	0	29	F	F	F
2001/2003	8	0	30	0	..	..	F	F	F
Saskatchewan									
1992/1994	43	13	72	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	46	17	74	56	21	84	F	F	F
2000/2002	42	15	68	44	14	74	F	F	F
2001/2003	48	18	75	F	F	F	F	F	F
Alberta									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	21	3	50	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	23	4	55	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	17	3	44	F	F	F	F	F	F
1998/2000	27	8	51	36	11	64	F	F	F
1999/2001	19	6	38	25	8	47	F	F	F
2000/2002	29	13	48	37	15	60	F	F	F
2001/2003	25	11	42	27	10	48	21	3	49
British Columbia									
1992/1994	37	23	52	36	21	52	F	F	F
1993/1995	28	16	42	27	14	43	32	8	61
1994/1996	33	20	46	31	17	46	37	11	66
1995/1997	37	23	51	34	19	51	43	16	71
1996/1998	32	18	47	30	15	47	38	12	67
1997/1999	16	5	32	16	4	35	F	F	F
1998/2000	27	13	44	23	8	42	F	F	F
1999/2001	42	25	58	38	20	56	F	F	F
2000/2002	41	27	55	36	20	52	54	25	80
2001/2003	46	32	60	42	27	57	59	29	83

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).