

Table 9-2

Hypopharynx — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Hypopharynx (ICD-O-3: C12.9, C13.0-C13.9) ³									
1992/1994									
Canada	26	22	30	24	20	28	32	23	41
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	25	20	31	24	18	30	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	27	19	35	F	F	F	F	F	F
1993/1995									
Canada	25	21	29	26	21	30	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	24	19	30	25	19	31	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	24	16	34	F	F	F	F	F	F
1994/1996									
Canada	24	20	28	26	21	30	18	10	27
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	23	18	29	24	18	30	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1995/1997									
Canada	28	23	32	29	24	34	23	14	33
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	26	20	32	26	19	32	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1996/1998									
Canada	26	22	31	26	21	31	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	24	19	30	22	16	28	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F

See footnotes at the end of the table.

Table 9-2 – continued

Hypopharynx — Age-standardized five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, Canada and selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
1997/1999									
Canada	27	22	31	26	22	31	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	26	20	32	24	18	31	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1998/2000									
Canada	25	21	29	24	19	28	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	23	17	29	21	15	27	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
1999/2001									
Canada	27	22	31	26	21	30	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	25	19	30	23	17	29	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	31	21	43	F	F	F	F	F	F
2000/2002									
Canada	28	23	32	26	22	31	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	F	F	F
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	27	21	33	25	19	31	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	F	F	F	F	F	F	F	F	F
2001/2003									
Canada	29	25	34	29	24	34	F	F	F
Newfoundland and Labrador	F	F	F	F	F	F	..	..	..
Prince Edward Island	F	F	F	F	F	F	F	F	F
Nova Scotia	F	F	F	F	F	F	F	F	F
New Brunswick	F	F	F	F	F	F	F	F	F
Ontario	31	24	37	31	23	38	F	F	F
Manitoba	F	F	F	F	F	F	F	F	F
Saskatchewan	F	F	F	F	F	F	F	F	F
Alberta	F	F	F	F	F	F	F	F	F
British Columbia	25	16	36	25	15	37	F	F	F

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1572).