

Table 9-3

Hypopharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Hypopharynx (ICD-O-3: C12.9, C13.0-C13.9) ³									
Newfoundland and Labrador									
1992/1994	19 ^E	5 ^E	42 ^E	23 ^E	6 ^E	49 ^E	F	F	F
1993/1995	28 ^E	9 ^E	54 ^E	28 ^E	9 ^E	54 ^E	F	F	F
1994/1996	19 ^E	3 ^E	47 ^E	19 ^E	3 ^E	47 ^E	F	F	F
1995/1997	43 ^E	13 ^E	74 ^E	43 ^E	13 ^E	74 ^E	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	..	..	..
Prince Edward Island									
1992/1994	F	F	F	F	F	F	F	F	F
1993/1995	F	F	F	F	F	F	F	F	F
1994/1996	F	F	F	F	F	F	F	F	F
1995/1997	F	F	F	F	F	F	F	F	F
1996/1998	F	F	F	F	F	F	F	F	F
1997/1999	F	F	F	F	F	F	F	F	F
1998/2000	F	F	F	F	F	F	F	F	F
1999/2001	F	F	F	F	F	F	F	F	F
2000/2002	F	F	F	F	F	F	F	F	F
2001/2003	F	F	F	F	F	F	F	F	F
Nova Scotia									
1992/1994	29	12	50	27	10	49	F	F	F
1993/1995	40	19	63	47	22	71	F	F	F
1994/1996	37	19	58	44	22	66	F	F	F
1995/1997	31	14	52	40	18	64	F	F	F
1996/1998	5	0	22	7	0	27	F	F	F
1997/1999	15	2	39	18	3	44	F	F	F
1998/2000	14	4	33	18	4	40	F	F	F
1999/2001	16	4	36	23	6	48	F	F	F
2000/2002	8	1	24	11	2	30	F	F	F
2001/2003	17	4	38	20	5	44	F	F	F
New Brunswick									
1992/1994	47	20	73	37	11	66	F	F	F
1993/1995	30	9	55	25	6	51	F	F	F
1994/1996	36	11	63	36	11	63	F	F	F
1995/1997	25	8	48	25	8	48	F	F	F
1996/1998	22	7	44	22	7	44	F	F	F
1997/1999	20	6	41	20	6	41	F	F	F
1998/2000	15	4	33	10	2	29	F	F	F
1999/2001	20	6	41	17	4	39	F	F	F
2000/2002	21	7	42	18	4	40	F	F	F
2001/2003	28	9	53	29	9	56	F	F	F
Ontario									
1992/1994	25	20	31	25	19	32	25	14	37
1993/1995	25	19	30	26	20	33	18	10	30
1994/1996	24	18	30	25	18	31	21	11	33
1995/1997	26	20	32	26	19	33	27	15	41
1996/1998	24	18	30	22	16	28	34	20	49
1997/1999	25	19	31	24	18	31	29	15	46
1998/2000	23	18	29	21	16	27	33	18	50
1999/2001	26	21	32	25	19	31	35	19	52
2000/2002	28	22	34	26	19	32	39	23	54
2001/2003	31	25	38	30	23	38	34	20	55

See footnotes at the end of the table.

Table 9-3 – continued

Hypopharynx — Five-year relative survival ratios, October 2011 CCR file, by sex, population aged 15 to 99, 3 years of cases, selected provinces^{1,2}

Primary site of cancer	3 years of cases								
	Both sexes			Males			Females		
	Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval		Relative survival ratio	95% confidence interval	
		Low	High		Low	High		Low	High
percent									
Manitoba									
1992/1994	26	12	42	21	9	38	F	F	F
1993/1995	27	13	44	27	13	45	F	F	F
1994/1996	17	6	32	20	7	38	F	F	F
1995/1997	29	15	46	36	17	56	F	F	F
1996/1998	33	14	56	42	15	71	F	F	F
1997/1999	25	10	44	27	10	49	F	F	F
1998/2000	12	2	34	14	2	37	F	F	F
1999/2001	22	8	41	26	9	47	F	F	F
2000/2002	32	15	52	32	13	55	F	F	F
2001/2003	39	21	58	37	16	59	44	14	74
Saskatchewan									
1992/1994	33	12	58	23	6	49	F	F	F
1993/1995	30	12	51	22	7	45	F	F	F
1994/1996	27	12	46	25	9	46	F	F	F
1995/1997	36	18	55	39	18	60	F	F	F
1996/1998	30	12	52	39	16	64	F	F	F
1997/1999	35	13	60	34	11	63	F	F	F
1998/2000	21	5	46	18	3	45	F	F	F
1999/2001	31	11	55	31	10	58	F	F	F
2000/2002	36	13	62	39	14	66	F	F	F
2001/2003	35	14	58	37	15	61	F	F	F
Alberta									
1992/1994	24	13	37	20	9	34	F	F	F
1993/1995	21	10	36	21	9	36	F	F	F
1994/1996	25	12	40	30	15	48	F	F	F
1995/1997	28	14	45	32	14	52	20	3	49
1996/1998	31	17	46	35	17	54	23	6	49
1997/1999	24	12	39	22	9	39	32	8	62
1998/2000	25	13	38	24	12	39	F	F	F
1999/2001	27	15	40	24	12	38	F	F	F
2000/2002	27	16	40	25	13	38	F	F	F
2001/2003	22	12	35	21	11	35	F	F	F
British Columbia									
1992/1994	26	18	35	27	17	38	25	12	41
1993/1995	25	16	35	26	16	37	21	7	42
1994/1996	24	15	34	26	16	37	16	3	40
1995/1997	26	17	37	27	17	39	18	3	45
1996/1998	29	18	40	29	18	42	27	8	52
1997/1999	29	19	40	32	20	44	19	5	42
1998/2000	33	23	44	32	21	45	36	13	62
1999/2001	34	23	45	36	24	49	26	8	49
2000/2002	34	23	46	32	20	45	42	15	69
2001/2003	28	18	40	28	18	40	29	7	58

1. Data from Quebec have been excluded because of issues in correctly ascertaining the vital status of cases.

2. In the years under study, there was a known under reporting of cancer cases in Newfoundland and Labrador. There is likely to be some overestimation of survival for this province as the survival of such "missed" cases is generally less favourable than that of cases in the registry population. As such, survival estimates from the province of Newfoundland and Labrador should be treated with caution.

3. Excluding morphology types M-9050 to M-9055; M-9140; M-9590 to M-9989.

Note(s): See "Data quality, concepts and methodology — Footnotes" section. Please note that CANSIM tables also provide the number of cases on which analyses were based, the number of deaths within the first five years of follow-up, as well as estimates of observed survival and their corresponding 95% confidence intervals. Survival estimates based on cases from individual years are available through CANSIM tables.

Source(s): Statistics Canada, Canadian Cancer Registry (CCR), Canadian Vital Statistics (Death Database) and life tables (CANSIM table 103-1574).