

## Article

### Health Fact Sheet

# Body composition of Canadian adults, 2009 to 2011

Health Statistics Division

September 2012



## How to obtain more information

For information about this product or the wide range of services and data available from Statistics Canada, visit our website, [www.statcan.gc.ca](http://www.statcan.gc.ca), email us at [infostats@statcan.gc.ca](mailto:infostats@statcan.gc.ca), or telephone us, Monday to Friday from 8:30 a.m. to 4:30 p.m., at the following numbers:

### Statistics Canada's National Contact Centre

Toll-free telephone (Canada and United States):

Inquiries line	1-800-263-1136
National telecommunications device for the hearing impaired	1-800-363-7629
Fax line	1-877-287-4369

Local or international calls:

Inquiries line	1-613-951-8116
Fax line	1-613-951-0581

### Depository Services Program

Inquiries line	1-800-635-7943
Fax line	1-800-565-7757

## To access this product

This product, Catalogue no. 82-625-X, is available free in electronic format. To obtain a single issue, visit our website, [www.statcan.gc.ca](http://www.statcan.gc.ca), and browse by "Key resource" > "Publications."

## Standards of service to the public

Statistics Canada is committed to serving its clients in a prompt, reliable and courteous manner. To this end, Statistics Canada has developed standards of service that its employees observe. To obtain a copy of these service standards, please contact Statistics Canada toll-free at 1-800-263-1136. The service standards are also published on [www.statcan.gc.ca](http://www.statcan.gc.ca) under "About us" > "The agency" > "Providing services to Canadians."

Published by authority of the Minister responsible for  
Statistics Canada

© Minister of Industry, 2012

All rights reserved. Use of this publication is governed by the  
Statistics Canada Open Licence Agreement ([http://www.  
statcan.gc.ca/reference/copyright-droit-auteur-eng.htm](http://www.statcan.gc.ca/reference/copyright-droit-auteur-eng.htm)).

Cette publication est aussi disponible en français.

## Note of appreciation

Canada owes the success of its statistical system to a long-standing partnership between Statistics Canada, the citizens of Canada, its businesses, governments and other institutions. Accurate and timely statistical information could not be produced without their continued co-operation and goodwill.

## Standard symbols

The following symbols are used in Statistics Canada publications:

.	not available for any reference period
..	not available for a specific reference period
...	not applicable
0	true zero or a value rounded to zero
0 <sup>s</sup>	value rounded to 0 (zero) where there is a meaningful distinction between true zero and the value that was rounded
P	preliminary
r	revised
X	suppressed to meet the confidentiality requirements of the <i>Statistics Act</i>
E	use with caution
F	too unreliable to be published
*	significantly different from reference category ( $p < 0.05$ )



## Body composition of Canadian adults, 2009 to 2011

In 2009 to 2011, 67% of Canadian men and 54% of Canadian women aged 18 to 79 were overweight or obese based on results from the Canadian Health Measures Survey.


Body composition is an important indicator of the health of individuals and populations. The body mass index (BMI) is used to identify weight-related health risks by classifying individuals as either underweight, normal weight, overweight or obese.<sup>1</sup>

Overweight and obesity are associated with an increased risk of numerous health problems including type 2 diabetes, hypertension, obstructive sleep apnea, osteoarthritis, many types of cancer (including breast, colorectal and pancreatic) and cardiovascular disease (coronary heart disease and stroke).<sup>1, 2</sup> Being underweight is also associated with health problems, including osteoporosis, under-nutrition, infertility and an increased risk of mortality.<sup>1, 3</sup>

Among Canadians aged 18 to 79, almost 2 out of every 3 (60%) adults aged 18 to 79 were overweight or obese while 2% were underweight, based on their measured BMI.

While 40% of Canadian men aged 18 to 79 were overweight and 27% were obese, 32% had a normal BMI (Chart 1). For women, 44% of Canadians aged 18 to 79 had a normal BMI, while 29% were overweight, 25% were obese and 3% were underweight (Chart 2).

**Chart 1 Distribution of males aged 18 to 79, by BMI category† and age group, Canada, 2009 to 2011\***



† BMI classification based on Canadian Guidelines for Body Weight Classification in Adults<sup>1</sup>

\* Data for the underweight category is too unreliable to be published (data with a coefficient of variation (CV) greater than 33.3%; suppressed due to extreme sampling variability).


Source: Canadian Health Measures Survey, 2009 to 2011. The CHMS collects health information on the Canadian household population aged 3 to 79.



For both adult men and women, those aged 18 to 39 were the most likely to have a normal BMI. Men and women in this age group were also significantly less likely to be overweight or obese than their counterparts aged 40 and over.

Men were significantly more likely to be overweight or obese than women. While approximately 3 out of 10 Canadian men had normal body weight, almost 7 out of 10 were overweight or obese. Among women, more than 4 out of 10 had a normal weight while more than 5 out of 10 were overweight or obese.

**Chart 2 Distribution of females aged 18 to 79, by BMI category† and age group, Canada, 2009 to 2011**



† BMI classification based on Canadian Guidelines for Body Weight Classification in Adults<sup>1</sup>

<sup>E</sup> Use with caution (data with a coefficient of variation (CV) from 16.6% to 33.3%)

<sup>F</sup> Too unreliable to be published (data with a coefficient of variation (CV) greater than 33.3%; suppressed due to extreme sampling variability)

Source: Canadian Health Measures Survey, 2009 to 2011. The CHMS collects health information on the Canadian household population aged 3 to 79.




## Waist circumference

As BMI does not provide an indication of fat distribution, waist circumference measurements are used to provide an estimation of abdominal obesity. Abdominal fat can result in increased risk of health problems as excess fat around the waist and upper body is associated with greater health risks than fat located more in the hip and thigh areas.<sup>1</sup>

Health Canada's health risk classification suggests that a waist circumference of greater than or equal to 102cm for men and greater than or equal to 88cm for women, can lead to an increased risk for developing health problems such as type 2 diabetes, coronary heart disease and hypertension.<sup>1</sup>

The average waist circumference for Canadian men aged 18 to 79 was 95.1 cm and for women it was 87.3 cm. Among Canadian males, 29% had a waist circumference associated with increased health risks. Canadian women however were significantly more likely to have a waist circumference above the cut-off, with 41% having a waist circumference associated with an increased health risk.

**Chart 3** Distribution of household population aged 18 to 79 with increased waist circumference risk,<sup>†</sup> by sex, Canada, 2009 to 2011



<sup>†</sup> Based on Canadian Guidelines for Body Weight Classification in Adults<sup>1</sup>

Source: Canadian Health Measures Survey, 2009 to 2011. The CHMS collects health information on the Canadian household population aged 3 to 79.

## About body mass index

Body mass index (BMI) is the ratio of person's weight in kilograms by their height squared in meters (kg/m<sup>2</sup>). BMI can be interpreted using the following classification table provide by Health Canada<sup>1</sup>:

BMI (kg/m <sup>2</sup> )	Classification	Risk of developing health problems
Less than 18.5	Underweight	Increased
18.5 to 24.9	Normal	Least
25.0 to 29.9	Overweight	Increased
30.0 to 34.9	Obese, Class I	High
35.0 to 39.9	Obese, Class II	Very high
40.0 and over	Obese, Class III	Extremely high

## References

1. *Canadian Guidelines for Body Weight Classification in Adults*. Ottawa: Health Canada; 2003.
2. *Obesity in Canada*. Ottawa: Public Health Agency of Canada, Canadian Institute for Health Information; 2011.
3. Orpana H, Berthelot JM, Kaplan MS, Feeny DH, McFarland B, Ross NA. BMI and mortality: results from a national longitudinal study of Canadian adults. *Obesity*. 2010;18(1):214-8.



# Canadian Health Measures Survey

For more information on the Canadian Health Measures Survey, please contact Statistics Canada's National Contact Centre (toll-free 1-800-263-1136; 613-951-8116; [infostats@statcan.gc.ca](mailto:infostats@statcan.gc.ca)).